

BE.
Third Semester Examination, May-2009
MATHEMATICS-III

Note : Attempt any five questions, selecting at least one question from each part.

Part-A

Q. 1. (a) Find the Fourier series for the function,

$$f(x) = \begin{cases} x & -1 < x \leq 0 \\ x+2 & 0 < x < 1 \end{cases}$$

where $f(x) = f(x+2)$

Ans.
$$f(x) = \begin{cases} x & -1 < x \leq 0 \\ x+2 & 0 < x < 1 \end{cases}$$

Here interval is $(-1, 1)$. Here $C = 1$

$$a_0 = \frac{1}{C} \int_{-C}^C f(x) dx = \frac{1}{1} \int_{-1}^1 f(x) dx$$

$$= \int_{-1}^0 x dx + \int_0^1 (x+2) dx$$

$$= \left[\frac{x^2}{2} \right]_{-1}^0 + \left[\frac{x^2}{2} + 2x \right]_0^1$$

$$= -\frac{1}{2} + \frac{1}{2} + 2 = 2$$

$$a_n = \frac{1}{C} \int_{-C}^C f(x) \cos \frac{n\pi x}{C} dx$$

$$= \frac{1}{1} \int_{-1}^1 f(x) \cos n\pi x dx$$

$$= \int_{-1}^0 x \cos n\pi x dx + \int_0^1 (x+2) \cos n\pi x dx$$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

$$= \left[x \left(\frac{\sin n\pi x}{n\pi} \right) - (1) \left(-\frac{\cos n\pi x}{n^2 \pi^2} \right) \right]_{-1}^0 + \left[(x+2) \frac{\sin n\pi x}{n\pi} + \frac{\cos n\pi x}{n^2 \pi^2} \right]_0^1$$

$$= \frac{1}{n^2 \pi^2} - \frac{(1)^0}{n^2 \pi^2} + \frac{(-1)^n}{n^2 \pi^2} - \frac{1}{n^2 \pi^2} = 0$$

$$b_n = \frac{1}{+1} \int_{-1}^1 f(x) \frac{\sin n\pi x}{1} dx$$

$$= \int_{-1}^0 x \sin n\pi x dx + \int_0^1 (x+2) \sin n\pi x dx$$

$$\Rightarrow \left[x \left(-\frac{\cos n\pi x}{n\pi} \right) - (1) \left(-\frac{\sin n\pi x}{n^2 \pi^2} \right) \right]_{-1}^0 + \left[(x+2) \left(-\frac{\cos n\pi x}{n\pi} \right) + (1) \frac{\sin n\pi x}{n^2 \pi^2} \right]_0^1$$

$$= \left(\frac{(-1)(-1)(-1)^n}{n\pi} \right) + \left(\frac{-3(-1)^n}{n\pi} + \frac{2}{n\pi} \right) = \frac{-4(-1)^n}{n\pi} + \frac{1}{n\pi}$$

$$f(x) = \frac{a_0}{2} + \sum a_n \cos \frac{n\pi x}{C} + \sum b_n \frac{\sin n\pi x}{C}$$

$$f(x) = 1 + \sum_{n=1}^{\infty} \left(\frac{-4(-1)^n}{n\pi} + \frac{2}{2\pi} \right) \frac{\sin n\pi x}{1}$$

$$f(x) = 1 + \frac{6}{\pi} \sin \pi x - \frac{2}{2\pi} \sin 2\pi x + \dots \infty$$

Q. 1. (b) Find the Fourier cosine and sine series of

$$f(x) = 1, 0 \leq x \leq 2.$$

Ans. $f(x) = 1 \quad 0 < x < 2$

(i) For the Half Range Sine Series :

$$1 = \sum b_n \sin \frac{n\pi x}{2} \quad (\text{since } l=2)$$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

$$b_n = \frac{2}{2} \int_1^2 1 \sin \frac{n\pi x}{2} dx$$

$$= \left[\frac{-\cos \frac{n\pi x}{2}}{\frac{n\pi}{2}} \right]_1^2 = -\frac{2}{n\pi} ((-1)^n - 1)$$

$$= \frac{4}{n\pi}, \text{ if } n \text{ is odd}$$

0, if n is even

$$f = \frac{4}{\pi} \sin \frac{\pi x}{2} + \frac{4}{3\pi} \sin \frac{3\pi x}{2} + \frac{4}{5\pi} \sin \frac{5\pi x}{2} + \dots$$

(ii) For Half Range Cosine Series :

$$f \approx \frac{a_0}{2} + \sum a_n \cos \frac{n\pi x}{2}$$

$$a_0 = \frac{2}{2} \int_0^2 1 dx = [x]_0^2 = 2$$

$$a_n = \frac{2}{2} \int_1^2 1 \cos \frac{n\pi x}{2} dx = \left[\frac{\sin \frac{n\pi x}{2}}{\frac{n\pi}{2}} \right]_1^2 = 0$$

$$\boxed{f(x) = 1 + 0}$$

Q. 2. (a) Find the Fourier transform of

$$f(x) = xe^{-ax^2}, a > 0.$$

Ans.

$$F[f(x)] = F[e^{-ax^2}]$$

$$\text{First to find } F[e^{-x^2}]$$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

Let

$$f(x) = e^{-x^2}$$

Then

$$\begin{aligned} F[f(x)] &= \int_{-\infty}^{\infty} f(x) e^{isx} dx \\ &= \int_{-\infty}^{\infty} e^{-x^2} e^{isx} dx \\ &= \int_{-\infty}^{\infty} e^{-\left(x^2 - isx\right)} dx = \int_{-\infty}^{\infty} e^{-\left[\left(x - \frac{is}{2}\right)^2 + \frac{s^2}{4}\right]} dx \\ &= e^{-s^2/4} \int_{-\infty}^{\infty} e^{-\left(x - \frac{is}{2}\right)^2} dx \\ &= e^{-s^2/4} \int_{-\infty}^{\infty} e^{-z^2} dz \quad \text{where } z = x - \frac{is}{2} \\ &= 2e^{-s^2/4} \int_0^{\infty} e^{-z^2} dz = 2e^{-s^2/4} \frac{\sqrt{\pi}}{2} \\ &= \sqrt{\pi} e^{-s^2/4} = F(s) \end{aligned}$$

Now,

$$e^{-ax^2} = e^{-\left(\sqrt{ax}\right)^2} = f(\sqrt{ax})$$

By change of scale property, we have

$$F[f\sqrt{ax}] = \frac{1}{\sqrt{a}} F\left(\frac{s}{\sqrt{a}}\right)$$

$$F\left[e^{-ax^2}\right] = \frac{1}{\sqrt{a}} \sqrt{\pi} e^{-\frac{1}{4}\left(\frac{s}{\sqrt{a}}\right)^2}$$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

$$= \sqrt{\frac{\pi}{a}} e^{-s^2/4a}$$

Thus,

$$\boxed{F[e^{-ax^2}] = \sqrt{\frac{\pi}{a}} e^{-s^2/4a}}$$

Q. 2. (b) State and prove time-shifting and frequency-shifting properties for Fourier transformation.

Ans. Shifting on time axis : If $F(s)$ is the complex Fourier transform of $f(t)$ and t_0 is any real number, then

$$F[f(t - t_0)] = e^{-ist_0} F(s)$$

$$\therefore F(f(t - t_0)) = \int_{-\infty}^{\infty} f(t - t_0) e^{ist} dt \quad \dots (1)$$

Putting $t - t_0 = T$, i.e. $t = t_0 + T$, we have $dt = dT$, when $t \rightarrow -\infty$, $T \rightarrow -\infty$ and when $t \rightarrow \infty$, $T \rightarrow \infty$.
From equation (1), we have

$$\begin{aligned} F[f(t - t_0)] &= \int_{-\infty}^{\infty} f(T) e^{is(t_0 + T)} dT \\ &= \int_{-\infty}^{\infty} f(T) e^{is t_0} e^{isT} dT \\ &= e^{is t_0} \int_{-\infty}^{\infty} f(T) e^{isT} dT \\ &= e^{is t_0} F(s) \end{aligned}$$

Shifting on frequency axis :

If $F(s)$ is the complex Fourier transform of $f(t)$ and s_0 is any real number then,

$$F[e^{is_0 t} f(t)] = F(s + s_0)$$

Proof : By definition we have

$$F(s) = F(f(t)) = \int_{-\infty}^{\infty} f(t) e^{ist} dt$$

$$F(e^{is_0 t} f(t)) = \int_{-\infty}^{\infty} e^{is_0 t} f(t) e^{ist} dt$$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

$$= \int_{-\infty}^{\infty} f(t) e^{i(s+s_0)t} dt = F(s+s_0).$$

Part-B

Q. 3. (a) If $f(z)$ is analytic function with constant modulus, then prove that $f(z)$ is constant.

Ans. Let $f(z) = u + iv$ be an analytic function.

Since $|f(z)| = \text{constant} = c$ (say) ($c \neq 0$)

$$\therefore |f(z)|^2 = u^2 + v^2 = c^2 \quad \dots(1)$$

Diff. (1) partially w.r.t x and y respectively, we have

$$2u \frac{\partial u}{\partial x} + 2v \frac{\partial v}{\partial x} = 0$$

$$\text{or} \quad \frac{u \partial u}{\partial x} + \frac{v \partial v}{\partial x} = 0 \quad \dots(2)$$

$$\text{and} \quad 2u \frac{\partial u}{\partial y} + 2v \frac{\partial v}{\partial y} = 0$$

$$\text{or} \quad \frac{u \partial u}{\partial y} + \frac{v \partial v}{\partial y} = 0 \quad \dots(3)$$

Using C-R equations,

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$$

$$\text{and} \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

Equation (3) becomes,

$$\therefore \frac{\partial v}{\partial x} u - \frac{\partial u}{\partial x} v = 0 \quad \dots(4)$$

Squaring and adding (2) and (4), we have

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

$$(u^2 + v^2) \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \right] = 0$$

$$C^2 \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 \right] = 0 \quad \because C \neq 0$$

$$\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial v}{\partial x} \right)^2 = 0$$

$$|f'(z)|^2 = 0 \quad \text{since } f'(z) = \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$$

$$f'(z) = 0 \Rightarrow f(z) = \text{constant}$$

Q. 3. (b) Show that the function

$$v(x, y) = \ln(x^2 + y^2) + x - 2y$$

is harmonic. Find its conjugate harmonic function $u(x, y)$ and the corresponding analytic function $f(z)$.

Aus.

$$v(x, y) = \ln(x^2 + y^2) + x - 2y$$

$$v_x = \frac{1}{x^2 + y^2} \cdot 2x + 1$$

$$v_{xx} = \frac{(x^2 + y^2)(2) - (2x)(2x)}{(x^2 + y^2)^2}$$

$$= \frac{2y^2 - 2x^2}{(x^2 + y^2)^2} = \frac{2(y^2 - x^2)}{(x^2 + y^2)^2}$$

$$v_y = \frac{2y}{x^2 + y^2} - 2$$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

$$v_{yy} = \frac{(x^2 + y^2)(2) - (2y)(2y)}{(x^2 + y^2)^2} = \frac{2x^2 - 2y^2}{(x^2 + y^2)^2}$$

Since $v_{xx} + v_{yy} = 0 \Rightarrow v$ is harmonic

Let u be harmonic conjugate of v the

$$du = u_x dx + u_y dy$$

$$= v_y dx - v_x dy$$

$$= \left(\frac{2y}{x^2 + y^2} - 2 \right) dx - \left(\frac{2x}{x^2 + y^2} + 1 \right) dy$$

$$u = \log \left| \frac{x+y}{x-y} \right| - 2x - \frac{2x}{2x} \log \frac{y+x}{y-y} - y$$

$$\log \frac{x+y}{x-y} - 2x - \log \frac{x+y}{x-y} - y$$

$$= -2x - y + c$$

$$f(z) = \left(\ln(x^2 + y^2)(x - 2y) \right) \hat{i} - (y + 2x)$$

$$= -(y + 2x) + \left[\ln(x^2 + y^2) + (x - 2y) \right] \hat{i}$$

$$\boxed{Z + iZ + 2\ln Z}$$

Q. 4. (a) Solve $\tan z = e^{i\alpha}$, where α is real.

Ans. $\tan z = e^{i\alpha} = \cos \alpha + i \sin \alpha$

As $\tan(x + iy) = \cos \alpha + i \sin \alpha$

$$\tan(x - iy) = \cos \alpha - i \sin \alpha$$

$$\tan 2\alpha = \tan(x + iy + x - iy)$$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

$$\begin{aligned}
 &= \frac{\tan(x+iy) + \tan(x-iy)}{1 - \tan(x+iy)\tan(x-iy)} \\
 &= \frac{(\cos\alpha + i\sin\alpha) + (\cos\alpha - i\sin\alpha)}{1 - (\cos\alpha + i\sin\alpha)(\cos\alpha - i\sin\alpha)} \\
 &= \frac{2\cos\alpha}{1 - (\cos^2\alpha + \sin^2\alpha)} = \frac{2\cos\alpha}{0}
 \end{aligned}$$

$$\tan 2x = \infty$$

Or

$$2x = \tan^{-1} \infty = 2n\pi + \frac{\pi}{2}$$

$$x = n\pi + \frac{\pi}{4} \quad \dots (i)$$

Again

$$2iy = (x+iy) - (x-iy)$$

$$\begin{aligned}
 \tan(2iy) &= \tan[(x+iy) - (x-iy)] \\
 &= \frac{\tan(x+iy) - \tan(x-iy)}{1 + \tan(x+iy)\tan(x-iy)} \\
 &= \frac{(\cos\alpha + i\sin\alpha) - (\cos\alpha - i\sin\alpha)}{1 + (\cos\alpha + i\sin\alpha)(\cos\alpha - i\sin\alpha)} \\
 &= \frac{2i\sin\alpha}{2} = i\sin\alpha
 \end{aligned}$$

Or

$$i \tanh 2y = i \sin\alpha$$

$$\tanh 2y = \sin\alpha$$

$$2y = \tanh^{-1} \sin\alpha$$

Using

$$\tanh^{-1} x = \frac{1}{2} \log \left(\frac{1+x}{1-x} \right)$$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

$$= \frac{1}{2} \log \left[\frac{1 + \sin \alpha}{1 - \sin \alpha} \right]$$

$$y = \frac{1}{4} \log \left[\frac{1 - \cos \left(\frac{\pi}{2} + \alpha \right)}{1 + \cos \left(\frac{\pi}{2} + \alpha \right)} \right]$$

$$= \frac{1}{4} \log \left[\frac{2 \sin^2 \left(\frac{\pi}{4} + \frac{\alpha}{2} \right)}{2 \cos^2 \left(\frac{\pi}{4} + \frac{\alpha}{2} \right)} \right]$$

$$y = \frac{1}{2} \log \tan \left(\frac{\pi}{4} + \frac{\alpha}{2} \right) \quad \dots(ii)$$

From equation (i) and (ii)

$$Z = x + iy = n\pi + \frac{\pi}{4} + \frac{1}{2} i \log \tan \left(\frac{\pi}{4} + \frac{\alpha}{2} \right).$$

Q. 4. (b) Evaluate $\oint_C \frac{dz}{z^2 + 9}$, where C is

(i) $|z + 3i| = 2$

(ii) $|z| = 5$

taken in counter-clockwise sense.

Ans. $\oint_C \frac{dz}{z^2 + 9}$

Here $f(z) = \frac{1}{z^2 + 9}$ has simple poles at $z = \pm 3i$

(i) $|z + 3i| = 2$

Only the pole $z = 3i$ lies inside the circle

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

Residue at $z = 3i$

$$\lim_{z \rightarrow 3i} (z - 3i) \frac{1}{(z^2 + 3i)(z - 3i)} = \frac{1}{6i} = -\frac{i}{6}$$

∴ By residue the,

$$\oint_C \frac{dz}{z^2 + 9} = 2\pi i \left(-\frac{i}{6} \right) = \frac{\pi}{3}$$

(ii) $|z| = 5$ is circle of centre $(0, 0)$ and radius 5

Both the poles $z = 3i$ and $z = -3i$ lies inside the circle.

Residue at $z = 3i$ is already calculate it comes $-\frac{i}{6}$.

Residue at $z = -3i$

$$\lim_{z \rightarrow -3i} (z + 3i) \frac{1}{(z + 3i)(z - 3i)} = \frac{1}{-6i} = \frac{i}{6}$$

By Residue theorem,

$$\begin{aligned} \oint_C \frac{dz}{z^2 + 9} &= 2\pi i \text{ [Sum of residue at both poles]} \\ &= 2\pi i \left[-\frac{i}{6} + \frac{i}{6} \right] = 0. \end{aligned}$$

Q. 5. (a) Explain $f(z) = \frac{7z-2}{z(z+1)(z-2)}$ as a Laurent series in the region $1 < |z+1| < 3$.

Ans. $\frac{7z-2}{z(z+1)(z-2)} \quad 1 < |z+1| < 3$

$$\frac{7z-2}{z(z+1)(z-2)} = \frac{A}{z} + \frac{B}{z+1} + \frac{C}{z-2} \quad \dots (1)$$

$$7z-2 = A(z+1)(z-2) + Bz(z-2) + Cz(z+1)$$

FOR MORE STUDY MATIRIEL LOG ON TO <http://studentsuvidha.in>

$$A = 1 \quad B = -2 \quad C = 2$$

Equation (1) reduces to,

$$\frac{1}{z} - \frac{3}{z+1} + \frac{2}{z-2} \quad \text{Since } 1 < |z+1| < 3$$

$$\frac{1}{z} - \frac{3}{z+1} + \frac{2}{-2\left(1 - \frac{z}{2}\right)} \quad 0 < z < 2$$

$$\frac{1}{z} - \frac{3}{z+1} + \frac{2}{-2\left(1 - \frac{z}{2}\right)^{-1}} \quad \left|\frac{z}{2}\right| < 1$$

$$= \frac{1}{z} - \frac{3}{z+1} - \left(1 + \frac{z}{2} + \left(\frac{z}{2}\right)^2 + \left(\frac{z}{2}\right)^3 + \dots \infty\right)$$

$$= \frac{1}{z} - 3(1+z)^{-1} - \left(1 + \frac{z}{2} + \left(\frac{z}{2}\right)^2 + \left(\frac{z}{2}\right)^3 + \dots \infty\right)$$

$$= \frac{1}{z} - 3(1+z)^{-1} + \left(1 + \frac{z}{2} + \left(\frac{z}{2}\right)^2 + \left(\frac{z}{2}\right)^3 + \dots \infty\right)$$

$$\frac{1}{z} - 3(1 - z + z^2 - z^3 + z^4 + \dots) - \left(1 + \frac{z}{2} + \left(\frac{z}{2}\right)^2 + \left(\frac{z}{2}\right)^3 + \dots \infty\right)$$

Q. 5. (b) Evaluate the integral $\int_{-\infty}^{\infty} \frac{\cos ax}{(x^2 + \alpha^2)(x^2 - \beta^2)} dx$.

Ans. $\int_{-\infty}^{\infty} \frac{\cos ax}{(x^2 + \alpha^2)(x^2 - \beta^2)} dx$

Consider $\int_C \frac{e^{iax}}{z^2 + \alpha^2}$; where C is closed curve consisting of upper half of the circle C_R whose equation

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

$|z| = R$ and a part of real axis from $-R$ to R .

$$\oint_C \frac{e^{inz}}{z^2 + a^2} = \int_{C_R} \frac{e^{iaz}}{z^2 + \alpha^2} + \int_{-R}^R \frac{e^{iaz}}{x^2 + \alpha^2}$$

Using Cauchy's residue theorem,

$$2\pi i [S_R] = \oint_C \frac{e^{inz}}{z^2 + a^2} = \int_{C_R} \frac{e^{iaz}}{z^2 + \alpha^2} + \int_{-R}^R \frac{e^{iaz}}{x^2 + \alpha^2}$$

$$f(z) = \frac{e^{iaz}}{z^2 + \alpha^2} \quad \dots(1)$$

$z = i\alpha$ is a simple pole inside 'C' $z = -i\alpha$ is a simple pole outside. Residue of $f(z)$ at $z = i\alpha$ is a simple pole.

$$= \lim_{z \rightarrow i\alpha} (z - i\alpha) f(z)$$


$$\text{Resi} = \lim_{z \rightarrow i\alpha} (z - i\alpha) \frac{e^{iaz}}{z^2 + \alpha^2}$$

$$\text{Resi} = \frac{e^{-m\theta}}{2i\alpha} \quad \dots(2)$$

C_R is semicircle

$$|z| = R, z = Re^{i\theta}$$

$$(dz) = R d\theta$$



FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

$$\theta: \theta \rightarrow 0$$

$$\left| \int_{C_R} \frac{e^{iaz}}{z^2 + \alpha^2} dz \right| < \int_{C_R} \frac{|e^{iaz}| |dz|}{|z|^2 - \alpha^2} \leq \int_0^\pi \frac{e^{-aR \sin \theta}}{R^2 - \alpha^2} R d\theta$$

Using Jordan in equality $0 \leq \theta \leq \frac{\pi}{2}$

$$\frac{2}{\pi} \leq \frac{\sin \theta}{\theta} \leq 1$$

Hence

$$\sin \theta \geq \frac{2\theta}{\pi}$$

$$e^{-aR \sin \theta} > e^{-aR \left(\frac{2\theta}{\pi} \right)}$$

$$\left| \int_{C_R} \frac{e^{iaz}}{z^2 + \alpha^2} dz \right| \leq \frac{R}{R^2 - \alpha^2} \int_0^\pi e^{-aR \sin \theta} d\theta$$

i.e.,

$$\leq \frac{R}{R^2 - \alpha^2} 2 \int_0^{\frac{\pi}{2}} e^{-\frac{2aR\theta}{\pi}} d\theta \leq \frac{2R}{R^2 - \alpha^2} \left[\frac{e^{-\frac{2aR\theta}{\pi}}}{-\frac{2aR}{\pi}} \right]_0^{\pi/2}$$

$$\leq \frac{\pi}{a(R^2 - \alpha^2)} [1 - e^{-aR}] \rightarrow 0 \text{ as } R \rightarrow \infty$$

and

$$\int_{-R}^R \frac{e^{iax}}{x^2 + \alpha^2} dx \rightarrow \int_{-\infty}^{\infty} \frac{e^{iax}}{x^2 + \alpha^2} dx \text{ as } R \rightarrow \infty$$

-(1)

Using equation (3) and (2) in (1)

$$2\pi i \frac{e^{-a\alpha}}{2\alpha i} = 0 \int_{-\infty}^{\infty} \frac{e^{iax}}{x^2 + \alpha^2} dx$$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

Hence,
$$\int_{-\infty}^{\infty} \frac{e^{iax}}{x^2 + a^2} dx = \frac{\pi e^{-a^2}}{a}$$

Equation real part,

$$\int_{-\infty}^{\infty} \frac{\cos ax}{x^2 + a^2} dx = \frac{\pi e^{-a^2}}{a}$$

Part-C

Q. 6. (a) Six cards are dealt from a well-shuffled deck of playing cards. Given that all six cards are black, find the probability that they are all of the same suit.

Ans. Probability that the six cards are drawn are of black colour,

$$\frac{{}^{26}C_6}{{}^{52}C_6}$$

Probability that six black cards drawn and of same suit,

$$\frac{4({}^{13}C_6)}{{}^{52}C_6} \times \frac{{}^{26}C_6}{{}^{52}C_6},$$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

Q. 7. (b) A cubical die is thrown 9000 times and a throw of 3 or 4 is observed 3240 times. Show that the die cannot be regarded as an unbiased one and find the extreme limits between which the probability of a throw of 3 or 4 lies.

Ans. Here, $n = 9000$

$P =$ Probability of success (i.e., getting 3 or 4 induce)

$$P = \frac{2}{6} = \frac{1}{3}, \quad q = 1 - \frac{1}{3} = \frac{2}{3}$$

$$p = \frac{X}{n} = \frac{3240}{9000} = 0.36$$

$$H_0 : \text{is unbiased, i.e., } P = \frac{1}{3}$$

$$H_1 : P \neq \frac{1}{3} \text{ (two tailed test)}$$

$$\text{The test statistic } Z = \frac{p - d}{\sqrt{\frac{PQ}{n}}}$$

$$= \frac{0.36 - 0.33}{\sqrt{\frac{1}{3} \times \frac{2}{3} \times \frac{1}{9000}}} = 0.03496$$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

$$|z| = 0.03494 < 1.96$$

Conclusion : Accept the hypothesis,

As $|z| < z_{\alpha}$, z_{α} is the tabulated value of z at 5% level of significance.

$\therefore H_0$ is accepted, we conclude that the die is unbiased.

To find 95% confidence, limits of the proportion it is given by $p \pm z_{\alpha} \sqrt{PQ/n}$

$$= 0.36 \pm 1.96 \sqrt{\left(\frac{2}{3} \times \frac{2}{3}\right) / 900}$$

$$= 0.36 \pm 0.01377$$

$$= (0.37377, 0.24623)$$

Q. 8. (a) Using Simplex method, maximize

$Z = 10x_1 + x_2 + 2x_3$ subject to $x_1 + x_2 - 2x_3 \leq 10$, $4x_1 + x_2 + x_3 \leq 20$, $x_1, x_2, x_3 \geq 0$.

Ans. $Z = 10x_1 + x_2 + 2x_3$

Introducing slack variable

$$x_1 + x_2 - 2x_3 + s_1 = 10$$

$$4x_1 + x_2 + x_3 + s_2 = 20$$

			C_j	10	1	2	0	0	Ratio
C_B	Basic	Sol b(X_R)	x_1	x_2	x_3	s_1	s_2		x_B / x
0	s_1	10	1	1	-2	1	0		$10/1 = 10$
$\leftarrow 0$	s_2	20	4	1	1	0	1		$20/4 = 5$
$Z=0$	x_3	$10 \uparrow$	1	1	2	0	0		
			$c_j = c_j - z_j$						
			c_j	10	1	2	0	0	$2 - \frac{10}{4}$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>

	z_3	s_1	5	0	$\frac{3}{4}$	$-\frac{9}{4}$	1	$-\frac{1}{4}$
10	x_1	5	1	$\frac{1}{4}$	$\frac{1}{4}$	0	$\frac{1}{4}$	
	z_3	50	0	$-\frac{2}{4}$	$-\frac{2}{4}$	0	0	

All z_j are zero or (-ve).

So, we obtain optimal solution.

$$x_1 = 5, x_2 = 0, x_3 = 0, \text{Max } z = 50$$

Which is required solution.

Q. 8. (b) Using Graphical method, solve the following LPP

Max $Z = 5x_1 + 3x_2$ subject to $3x_1 + 5x_2 \leq 15$, $5x_1 + 2x_2 \leq 10$, $x_1, x_2 \geq 0$.

Ams. **Max. $Z = 5x_1 + 3x_2$**

Subject to $3x_1 + 5x_2 \leq 15$

$5x_1 + 2x_2 \leq 10$

Corresponding equality is,

$$3x_1 + 5x_2 = 15$$

x_1	0	5
x_2	3	0


$$5x_1 + 2x_2 = 10$$

x_1	0	2
x_2	5	0

Corner points are A (2, 0), B (5, 0), C $\left(\frac{20}{19}, \frac{45}{19}\right)$

$$\text{Max } z = 5x_1 + 3x_2$$

FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>



At $A(2, 0) = 10$

$B(5, 0) = 15$

$C\left(\frac{25}{19}, \frac{45}{19}\right) = \frac{235}{19}$

Max value,

$\text{Max } z = \frac{235}{19}$

At $x_1 = \frac{20}{19}, x_2 = \frac{45}{19}$

S
FOR MORE STUDY MATERIAL LOG ON TO <http://studentsuvidha.in>