

B.TECH.
FIRST SEMESTER THEORY 2010-11

EAS-103

MATHEMATICS-I

Time: 3 Hours

Total Marks: 100

Note: Q. No. 1 is compulsory and carries 5 marks. Attempt one question from each unit, symbols have their usual meaning.

SECTION A

Q. 1. All parts of this question are compulsory:

(2 × 10 = 20)

(a) If $u = f\left(\frac{y}{x}\right)$ then

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \dots$$

Ans: 0

Justification: u is a homogeneous function of degree 0,

\therefore By Euler's theorem,

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = 0u = 0$$

(b) The curve $x^{2/3} + y^{2/3} = a^{2/3}$

is symmetrical about.....

Ans: Both x axis and y axis

Justification: Replacing x by $-x$ eqn remains unchanged, therefore curve is symmetrical about y axis. Replacing y by $-y$ eqn remains unchanged, therefore curve is symmetrical about x axis.

Indicate True or False of the following statements:

(c) (i) Two functions u and v are functionally dependent if their Jacobian with respect to x and y is zero. (True/False)

(ii) If $f(x, y) = 1 - x^2y^2$, then stationary point is $(0, 0)$. (True/False)

Ans: (c) (i) True (ii) True

Justification: $\frac{\partial f}{\partial x} = -2xy^2, \frac{\partial f}{\partial y} = -2x^2y$

$\frac{\partial f}{\partial x} = 0, \frac{\partial f}{\partial y} = 0$ gives $(x, y) = (0, 0)$, which is stationary point.

(d) (i) The minimum value of $f(x, y) = x^2 + y^2$ is zero. (True/False)

(ii) If u, v are functions of r, s are themselves function of x, y then $\frac{\partial(u, v)}{\partial(x, y)} = \frac{\partial(u, v)}{\partial(r, s)} \times \frac{\partial(r, s)}{\partial(x, y)}$. (True/False)

Ans: (d) (i) True (ii) False.

Pick the correct answer of the choices given below:

(e) The eigen values of $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$ are

(a) 0, 0, 0 (b) 0, 0, 1 (c) 0, 0, 3 (d) 1, 1, 1

Ans: (e) (c) 0, 0, 3

Justification: For eigen values $|A - \lambda I| = 0$

$$\Rightarrow \begin{vmatrix} 1-\lambda & 1 & 1 \\ 1 & 1-\lambda & 1 \\ 1 & 1 & 1-\lambda \end{vmatrix} = 0$$

$$\Rightarrow \begin{vmatrix} 3-\lambda & 1 & 1 \\ 3-\lambda & 1-\lambda & 1 \\ 3-\lambda & 1 & 1-\lambda \end{vmatrix} = 0$$

Applying $C_1 \Rightarrow C_1 + C_2 + C_3$

$$\Rightarrow (3-\lambda) \begin{vmatrix} 1 & 1 & 1 \\ 1 & 1-\lambda & 1 \\ 1 & 1 & 1-\lambda \end{vmatrix} = 0$$

$$\Rightarrow \lambda^2 (3-\lambda) = 0$$

$$\Rightarrow \lambda = 0, 0, 3$$

(f) The rank of the matrix $\begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$ is

(a) 0 (b) 1 (c) 2 (d) 3

Ans: (f) (c) 2

Justification: Given matrix = $\begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & -2 & -1 \\ 0 & 0 & 0 \end{bmatrix} \quad \begin{array}{l} \text{By } R_2 \rightarrow R_2 - R_1 \\ R_3 \rightarrow R_3 - R_1 \end{array}$$

\therefore Rank = No. of non zero rows in echelon form
= 2

(g) $\frac{\beta(m+1, n)}{\beta(m, n)}$ is equal to

(a) $\frac{m}{n}$ (b) $\frac{m+1}{n}$ (c) $\frac{m-1}{n}$ (d) $\frac{m}{m+n}$

Ans: (g) (d) $\frac{m}{m+n}$

Justification: $\frac{\beta(m+1, n)}{\beta(m, n)} = \frac{\frac{\Gamma(m+1)\Gamma(n)}{\Gamma(m+n+1)}}{\frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)}}$

$$\therefore \beta(m, n) = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)}$$

$$= \frac{\frac{\Gamma(m+1)\Gamma(n)\Gamma(m+n)}{\Gamma(m+n+1)}}{\frac{\Gamma(m)\Gamma(n)\Gamma(m+n)}{\Gamma(m+n)}} = \frac{m\Gamma(m)\Gamma(n)\Gamma(m+n)}{\Gamma(m)\Gamma(n)\Gamma(m+n)\Gamma(m+n)} = \frac{m}{m+n}$$

as $\Gamma(r+1) = r\Gamma(r) = \frac{m}{m+n}$

(h) The value of the integral $\int_0^{\infty} e^{-x^2} dx$ is

(a) $\frac{2}{\sqrt{\pi}}$ (b) $\frac{\sqrt{\pi}}{2}$ (c) $\frac{\pi}{2}$ (d) $\frac{2}{\pi}$

Ans. (b) $\frac{\sqrt{\pi}}{2}$

Fill up the blanks with the correct answer:

(i) The Gauss divergence theorem relates certain surface integrals to—

(volume integrals/line integrals)

Ans: Volume integrals

(j) The vector field $\vec{F} = x\hat{i} - y\hat{j}$ is divergence free—

(but not irrotational/and irrotational)

Ans: and irrotational.

Justification: $\nabla \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x & -y & 0 \end{vmatrix} = 0$

$\Rightarrow \vec{F}$ is irrotational.

SECTION B

Q. 2. Attempt any three parts of the following:

(10 × 3 = 30)

(a) If $y = \sin(a \sin^{-1} x)$. Find $(y_n)_{n=0}$.

Ans. $\because y = \sin(a \sin^{-1} x)$... (1)

$$\Rightarrow \frac{dy}{dx} = \cos(a \sin^{-1} x) \frac{a}{\sqrt{1-x^2}}$$

$$\Rightarrow y_1 = \frac{a}{\sqrt{1-x^2}} \cos(a \sin^{-1} x) \quad \dots (2)$$

$$\Rightarrow \sqrt{1-x^2} y_1 = a \cos(a \sin^{-1} x)$$

$$\Rightarrow (1-x^2) y_1^2 = a^2 \cos^2(a \sin^{-1} x)$$

$$= a^2 [1 - \sin^2(a \sin^{-1} x)]$$

$$\Rightarrow (1-x^2) y_1^2 = a^2 [(1-y^2)] \quad [\because y = \sin(a \sin^{-1} x)]$$

Differentiating it again, we get,

$$(1-x^2) 2y_1 y_2 - 2xy_1^2 = a^2 (-2yy_1) \\ \Rightarrow (1-x^2)y_2 - xy_1 + a^2 y = 0 \quad \dots(3)$$

Differentiating above expression n times by Leibnitz's theorem, we get,

$$[(1-x^2)y_{n+2} + {}^nC_1(-2x)y_{n+1} + {}^nC_2(-2)y_n] \\ - [x y_{n+1} + n C_1(1)y_n] + a^2 y_n = 0 \\ \Rightarrow (1-x^2)y_{n+2} + n(-2x)y_{n+1} + \frac{n(n-1)}{2}(-2)y_n \\ - xy_{n+1} - n(1)y_n + a^2 y_n = 0 \\ \left[\because {}^nC_1 = n, {}^nC_2 = \frac{n(n-1)}{2} \right] \\ \Rightarrow (1-x^2)y_{n+2} - (2n+1)x y_{n+1} \\ + (a^2 - n^2 + n - n)y_n = 0 \\ \Rightarrow (1-x^2)y_{n+2} - (2n+1)x y_{n+1} \\ + (a^2 - n^2)y_n = 0 \quad \dots(4)$$

Putting $x = 0$ in (1),

$$y(0) = \sin(a \sin^{-1} 0) = 0$$

Putting $x = 0$ in (2),

$$y_1(0) = \frac{a}{\sqrt{1-0}} \cos(a \sin^{-1} 0) \\ \Rightarrow y_1(0) = a \cos 0 \\ \Rightarrow y_1(0) = a [\because \cos 0 = 1] \\ \text{Putting } x = 0 \text{ in (3),} \\ (1-0^2)y_2(0) - 0 y_1(0) + a^2 y(0) = 0 \\ y_2(0) + a^2(0) = 0 [\because y(0) = 0] \\ \Rightarrow y_2(0) = 0$$

Thus,

$$y_1(0) = a \text{ and } y_2(0) = 0$$

Again, putting $x = 0$ in (4), we get,

$$(1-0^2)y_{n+2}(0) - (2n+1)(0) + (a^2 - n^2)y_n(0) = 0 \\ \Rightarrow y_{n+2}(0) = (n^2 - a^2)y_n(0) \quad \dots(5)$$

Now, two cases arise.

Case I: When n is even

$$\text{Putting } n = 2, (5) \text{ gives } y_4(0) = (2^2 - a^2)y_2(0) \\ = 0 [\because y_2(0) = 0]$$

$$\text{Putting } n = 4, (5) \text{ gives } y_6(0) = (4^2 - a^2)y_4(0) \\ = 0 [\because y_4(0) = 0]$$

$$\text{Putting } n = 6, (5) \text{ gives } y_8(0) = (6^2 - a^2)y_6(0) \\ = 0 [\because y_6(0) = 0]$$

Thus, $y_n(0) = 0$ if n is even.

Case II. When n is odd

$$\text{Putting } n = 1, (5) \text{ gives } y_3(0) = (1^2 - a^2)y_1(0)$$

$$\Rightarrow y_3(0) = (1^2 - a^2)a [\because y_1(0) = a]$$

$$\text{Putting } n = 3, (5) \text{ gives } y_5(0) = (3^2 - a^2)y_3(0)$$

$$\Rightarrow y_5(0) = (1^2 - a^2)(3^2 - a^2)a$$

$$\text{Putting } n = 5, (5) \text{ gives } y_7(0) = (5^2 - a^2)y_5(0)$$

$$\Rightarrow y_7(0) = (1^2 - a^2)(3^2 - a^2)(5^2 - a^2)a$$

$$\Rightarrow y_n(0) = (1^2 - a^2)(3^2 - a^2)(5^2 - a^2) \dots [(n-2)^2 - a^2]a$$

if n is odd

$$\text{Thus, } y_n(0) = \begin{cases} 0 & \text{if } n \text{ is even} \\ (1^2 - a^2)(3^2 - a^2) \dots [(n-2)^2 - a^2]a & \text{if } n \text{ is odd and } n \neq 1 \end{cases}$$

**

(b) If u, v, w are the roots of the equation.

$$(x-a)^3(x-b)^3 + (x-c)^3 = 0,$$

then find $\frac{\partial(u, v, w)}{\partial(a, b, c)}$.

Ans: Given equation is

$$(x-a)^3 + (x-b)^3 + (x-c)^3 = 0 \quad \dots(1)$$

Now,

$$(x-a)^3 = x^3 - a^3 - 3x^2a + 3xa^2$$

$$[\because (a-b)^3 = a^3 - b^3 - 3a^2b + 3ab^2]$$

$$\Rightarrow (x-a)^3 = x^3 - 3ax^2 + 3a^2x - a^3$$

Similarly,

$$(x-b)^3 = x^3 - 3bx^2 + 3b^2x - b^3$$

$$(x-c)^3 = x^3 - 3cx^2 + 3c^2x - c^3$$

Putting these values in (1), we get,

$$x^3 - 3ax^2 + 3a^2x - a^3 + x^3 - 3bx^2 + 3b^2x - b^3 \\ + x^3 - 3cx^2 + 3c^2x - c^3 = 0 \\ \Rightarrow 3x^3 - 3(a+b+c)x^2 + 3(a^2+b^2+c^2)x \\ - (a^3+b^3+c^3) = 0 \quad \dots(2)$$

Now, we know that if α, β and γ are the roots of eqn $ax^3 + bx^2 + cx + d = 0$, then

$$\alpha + \beta + \gamma = \frac{-b}{a}$$

$$\alpha\beta + \beta\gamma + \gamma\alpha = \frac{c}{a}$$

$$\alpha\beta\gamma = -\frac{d}{a}$$

Here, $\alpha = u, \beta = v, \gamma = w$

Therefore,

$$u + v + w = -\frac{3(a+b+c)}{-3}$$

$$\Rightarrow u + v + w = a + b + c$$

$$\Rightarrow u + v + w - a - b - c = 0 \quad \dots(3)$$

Also,

$$uv + vw + wu = \frac{3(a^2 + b^2 + c^2)}{3}$$

$$\Rightarrow uv + vw + wu = a^2 + b^2 + c^2$$

$$\Rightarrow uv + vw + wu - a^2 - b^2 - c^2 = 0 \quad \dots(4)$$

Similarly,

$$\Rightarrow uvw = \frac{-(a^3 + b^3 + c^3)}{3}$$

$$\Rightarrow uvw - \frac{(a^3 + b^3 + c^3)}{3} = 0 \quad \dots(5)$$

From (3), (4) and (5)

$$\text{Let } f_1 = u + v + w - a - b - c$$

$$f_2 = uv + vw + wu - a^2 - b^2 - c^2$$

$$f_3 = uvw - \frac{1}{3}(a^3 + b^3 + c^3)$$

We know that,

$$\frac{\partial(u, v, w)}{\partial(a, b, c)} = (-1)^3 \frac{\partial(f_1, f_2, f_3)}{\partial(u, v, w)} \quad \dots(6)$$

Now,

$$\frac{\partial(f_1, f_2, f_3)}{\partial(a, b, c)} = \begin{vmatrix} \frac{\partial f_1}{\partial a} & \frac{\partial f_1}{\partial b} & \frac{\partial f_1}{\partial c} \\ \frac{\partial f_2}{\partial a} & \frac{\partial f_2}{\partial b} & \frac{\partial f_2}{\partial c} \\ \frac{\partial f_3}{\partial a} & \frac{\partial f_3}{\partial b} & \frac{\partial f_3}{\partial c} \end{vmatrix}$$

$$= \begin{vmatrix} -1 & -1 & -1 \\ -2a & -2b & -2c \\ -a^2 & -b^2 & -c^2 \end{vmatrix}$$

$$= (-1)(-2)(-1) \begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{vmatrix}$$

$$\left[\begin{array}{l} \text{Taking } -1 \text{ from } R_1, \\ -R \text{ from } R_2 \text{ and} \\ -\text{from } R_3 \text{ common} \end{array} \right]$$

$$= -2 \begin{vmatrix} 1 & 0 & 0 \\ a & b-a & c-a \\ a^2 & b^2-a^2 & c^2-a^2 \end{vmatrix} \left[\begin{array}{l} \text{By } C_2 \rightarrow C_2 - C_1 \\ \text{and } C_3 \rightarrow C_3 - C_1 \end{array} \right]$$

$$= -2 \begin{vmatrix} 1 & 0 & 0 \\ a & b-a & c-a \\ a^2 & (b-a)(b+a) & (c-a)(c+a) \end{vmatrix}$$

$$= -2(b-a)(c-a) \begin{vmatrix} 1 & 0 & 0 \\ a & 1 & 1 \\ a^2 & b+a & c+a \end{vmatrix}$$

Taking $(b-a)$ from C_2 and $(C-a)$ From C_3 common

$$= -2(b-a)(c-a)(c+a-b-a)$$

$$= -2(b-a)(c-a)(c-b)$$

$$= -2(a-b)(b-c)(c-a)$$

$$\Rightarrow \frac{\partial(f_1, f_2, f_3)}{\partial(a, b, c)} = -2(a-b)(b-c)(c-a)$$

Now,

$$\frac{\partial(f_1, f_2, f_3)}{\partial(u, v, w)} = \begin{vmatrix} \frac{\partial f_1}{\partial u} & \frac{\partial f_1}{\partial v} & \frac{\partial f_1}{\partial w} \\ \frac{\partial f_2}{\partial u} & \frac{\partial f_2}{\partial v} & \frac{\partial f_2}{\partial w} \\ \frac{\partial f_3}{\partial u} & \frac{\partial f_3}{\partial v} & \frac{\partial f_3}{\partial w} \end{vmatrix}$$

$$= \begin{vmatrix} 1 & 1 & 1 \\ v+w & v+w & u+v \\ vw & uw & uv \end{vmatrix}$$

$$= \begin{vmatrix} 1 & 0 & 0 \\ v+w & u-v & u-w \\ vw & w(u-v) & v(u-w) \end{vmatrix} \quad \begin{array}{l} \text{by } c_2 \rightarrow c_2 - c_1 \\ c_3 \rightarrow c_3 - c_1 \end{array}$$

$$= (u-v)(u-w) \begin{vmatrix} 1 & 0 & 0 \\ v+w & 1 & 1 \\ vw & w & v \end{vmatrix}$$

Taking $(u-v)$ from c_2 and $(u-w)$ from c_3 common

$$= (u-v)(u-w)(v-w)$$

$$\Rightarrow \frac{\partial(f_1, f_2, f_3)}{\partial(u, v, w)} = -(u-v)(v-w)(w-v)$$

Putting $\frac{\partial(f_1, f_2, f_3)}{\partial(a, b, c)}$ and $\frac{\partial(f_1, f_2, f_3)}{\partial(u, v, w)}$ in (6),

we get,

$$\frac{\partial(u, v, w)}{\partial(a, b, c)} = (-1)^3 \frac{2(a-b)(b-c)(c-a)}{-(u-v)(v-w)(w-u)}$$

$$\Rightarrow \frac{\partial(u, v, w)}{\partial(a, b, c)} = - \frac{2(a-b)(b-c)(c-a)}{(u-v)(v-w)(w-u)} \quad \text{Ans.}$$

(c) Find the eigen values and eigen vectors of the matrix

$$A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$$

$$\text{Ans. Given matrix } A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$$

characteristic equation of A is

$$|A - \lambda I| = 0$$

$$\Rightarrow \begin{vmatrix} 8-\lambda & -6 & 2 \\ -6 & 7-\lambda & -4 \\ 2 & -4 & 3-\lambda \end{vmatrix} = 0$$

$$\Rightarrow (8-\lambda)[(7-\lambda)(3-\lambda)-16] + 6[-6(3-\lambda)+8] + 2[24-2(7-\lambda)] = 0$$

$$\Rightarrow (8-\lambda)[21-7\lambda-3\lambda+\lambda^2-16] + 6[-18+6\lambda+8] + 2[24-14+2\lambda] = 0$$

$$\Rightarrow (8-\lambda)(\lambda^2-10\lambda+5) + 6(6\lambda-10) + 2(2\lambda+10) = 0$$

$$\Rightarrow 8\lambda^2 - 80\lambda + 40 - \lambda^3 + 10\lambda^2 - 5\lambda + 36\lambda - 60 + 4\lambda + 20 = 0$$

$$\Rightarrow -\lambda^3 + 18\lambda^2 - 45\lambda = 0$$

$$\Rightarrow -\lambda(\lambda^2 - 18\lambda + 45) = 0$$

$$\Rightarrow \lambda(\lambda^2 - 15\lambda - 3\lambda + 45) = 0$$

$$\Rightarrow \lambda(\lambda-15)-3(\lambda-15)] = 0$$

$$\Rightarrow \lambda(\lambda-3)(\lambda-15) = 0$$

$$\Rightarrow \lambda = 0, 3, 15, \text{ which are eigen values of A.}$$

$$\text{Let } x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \text{ be an eigen vector of A}$$

corresponding to eigen value λ , then $(A - \lambda I)x = 0$

$$\begin{bmatrix} 8-\lambda & -6 & 2 \\ -6 & 7-\lambda & -4 \\ 2 & -4 & 3-\lambda \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \quad \dots(1)$$

For $\lambda = 0$, (1) gives

$$\begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow 8x_1 - 6x_2 + 2x_3 = 0$$

$$-6x_1 - 7x_2 + 4x_3 = 0$$

$$2x_1 - (-4x_2) + 3x_3 = 0$$

From first two equations,

$$\begin{bmatrix} x_1 \\ -6 & 2 \\ 7 & -4 \end{bmatrix} = \begin{bmatrix} x_2 \\ 8 & 2 \\ -6 & -4 \end{bmatrix} = \begin{bmatrix} x_3 \\ 8 & -6 \\ -6 & 7 \end{bmatrix}$$

$$\Rightarrow \frac{x_1}{10} = \frac{x_2}{20} = \frac{x_3}{20}$$

$$\Rightarrow \frac{x_1}{1} = \frac{x_2}{2} = \frac{x_3}{2}$$

$$\Rightarrow \text{Eigen vector for eigen value 0 is } \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$$

Now, for $\lambda = 3$ (1) gives,

$$\begin{bmatrix} 5 & -6 & 2 \\ -6 & 4 & -4 \\ 2 & -4 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow 5x_1 - 6x_2 + 2x_3 = 0$$

$$-6x_1 + 4x_2 - 4x_3 = 0$$

$$2x_1 - 4x_2 + 0x_3 = 0$$

From first two equations,

$$\begin{bmatrix} x_1 \\ -6 & 2 \\ 4 & -4 \end{bmatrix} = \begin{bmatrix} x_2 \\ 5 & 2 \\ -6 & -4 \end{bmatrix} = \begin{bmatrix} x_3 \\ 5 & -6 \\ -6 & 4 \end{bmatrix}$$

$$\Rightarrow \frac{x_1}{16} = \frac{x_2}{8} = \frac{x_3}{-16}$$

$$\Rightarrow \frac{x_1}{2} = \frac{x_2}{+1} = \frac{x_3}{-2}$$

$$\Rightarrow \text{Eigen vector for eigen value 3 is } \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix}$$

Again, for $\lambda = 15$ (1) gives,

$$\begin{bmatrix} -7 & -6 & 2 \\ -6 & -8 & -4 \\ 2 & -4 & -12 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow -7x_1 - 6x_2 + 2x_3 = 0$$

$$-6x_1 - 8x_2 - 4x_3 = 0$$

$$2x_1 - 4x_2 - 12x_3 = 0$$

From first two equations,

$$\begin{bmatrix} x_1 \\ -6 & 2 \\ -8 & -4 \end{bmatrix} = \begin{bmatrix} x_2 \\ -7 & 2 \\ -6 & -4 \end{bmatrix} = \begin{bmatrix} x_3 \\ -7 & -6 \\ -6 & -8 \end{bmatrix}$$

$$\Rightarrow \frac{x_1}{40} = \frac{x_2}{-40} = \frac{x_3}{20}$$

$$\Rightarrow \frac{x_1}{2} = \frac{x_2}{-2} = \frac{x_3}{1}$$

$$\Rightarrow \text{Eigen vector for eigen value 15 is } \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix}$$

Thus, eigen values of A are 0, 3 and 15 corresponding eigen vectors are

$$\begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix} \text{ and } \begin{bmatrix} 2 \\ -2 \\ 1 \end{bmatrix} \text{ respectively}$$


(d) Change the order of integration in

$$I = \int_0^2 \int_{x^2/4}^{3-x} xy dy dx$$

and hence evaluate it.

$$\text{Ans: Given integral } I = \int_{x=0}^2 \int_{y=\frac{x^2}{4}}^{3-x} xy dy dx$$

The region of integration is shown in the figure.



By changing the order of integration,

$$I = \int_{R_1} xy dx dy + \int_{R_2} xy dx dy$$

For R_1 , Limits of x are from 0 to $\sqrt{4y}$

Limits of y are from 0 to 1

For R_2 ,

Limits of x are from 0 to $3-y$

Limits of y are from 1 to 3.

$$\text{Thus } I = \int_{y=0}^1 \int_{x=0}^{\sqrt{4y}} xy dx dy + \int_{y=1}^3 \int_{x=0}^{3-y} xy dx dy$$

$$= \int_{y=0}^1 \left[\frac{x^2 y}{2} \right]_{x=0}^{\sqrt{4y}} dy + \int_{y=1}^3 \left[\frac{x^2 y}{2} \right]_{x=0}^{3-y} dy$$

$$= \frac{1}{2} \int_{y=0}^1 4y(y) dy + \frac{1}{2} \int_{y=1}^3 (3-y)^2 y dy$$

$$= 2 \int_{y=0}^1 y^2 dy + \frac{1}{2} \int_{y=1}^3 (9 + y^2 - 6y) y dy$$

$$= 2 \left[\frac{y^3}{3} \right]_{y=0}^1 + \frac{1}{2} \int_{y=1}^3 (y^3 - 6y^2 + 9y) dy$$

$$= 2 \left(\frac{1}{3} \right) + \frac{1}{2} \left[\frac{y^4}{4} - 6 \frac{y^3}{3} + \frac{9y^2}{2} \right]_{y=1}^3$$

$$= \frac{2}{3} + \frac{1}{2} \left[\frac{81}{4} - 54 + \frac{81}{2} - \frac{1}{4} + \frac{6}{3} - \frac{9}{2} \right]$$

$$= \frac{2}{3} + \frac{1}{2} \left[\left(\frac{81}{4} - \frac{1}{4} \right) + \left(\frac{81}{2} - \frac{9}{2} \right) - 54 + 2 \right]$$

$$= \frac{2}{3} + \frac{1}{2} [20 + 36 - 52] = \frac{2}{3} + \frac{1}{2} (4)$$

$$= \frac{8}{3} \text{ Ans.}$$

(c) Find the volume enclosed between the two surfaces $Z = 8 - x^2 - y^2$ and $Z = x^2 + 3y^2$.

Ans : Required volume = $\iiint_v dx dy dz$, where v is the given region.

For v ,

Limits of z are from $x^2 + 3y^2$ to $8 - x^2 - y^2$. For limits of y , we equate z from given surfaces.

$$\Rightarrow 8 - x^2 - y^2 = x^2 + 3y^2$$

$$\Rightarrow 2x^2 + 4y^2 = 8$$

$$\Rightarrow x^2 + 2y^2 = 4$$

Thus, limits of y are from $-\sqrt{\frac{4-x^2}{2}}$ to

$$+\sqrt{\frac{4-x^2}{2}}.$$

Limits of x are from -2 to $+2$.

\therefore Required volume

$$= \int_{x=-2}^2 \int_{y=-\sqrt{\frac{4-x^2}{2}}}^{\sqrt{\frac{4-x^2}{2}}} \int_{z=x^2+3y^2}^{8-x^2-y^2} dz dy dx$$

$$= \int_{x=-2}^2 \int_{y=-\sqrt{\frac{4-x^2}{2}}}^{\sqrt{\frac{4-x^2}{2}}} (8 - x^2 - y^2 - x^2 - 3y^2) dy dx$$

$$= \int_{x=-2}^2 \int_{y=-\sqrt{\frac{4-x^2}{2}}}^{\sqrt{\frac{4-x^2}{2}}} (8 - 2x^2 - 4y^2) dy dx$$

Putting $x = 2r \cos \theta$, $y = \sqrt{2} r \sin \theta$

$$dx dy = 2 \sqrt{2} r dr d\theta$$

r varies from 0 to 1

θ varies from 0 to 2π .

Required volume

$$= \int_0^{2\pi} \int_{r=0}^1 [8 - 2(4r^2)] 2\sqrt{2} r dr d\theta$$

$$= \int_0^{2\pi} \int_{r=0}^1 8(1-r^2) 2\sqrt{2} r dr d\theta$$

$$= 16\sqrt{2} \int_0^{2\pi} \left[\frac{r^2}{2} - \frac{r^4}{4} \right]_{r=0}^1 d\theta$$

$$= 16\sqrt{2} \int_0^{2\pi} \frac{1}{4} d\theta$$

$$= 4\sqrt{2} (2\pi)$$

$$= 8\sqrt{2} \pi \text{ Ans.}$$

SECTION C

Attempt any two parts from each equation. All questions are compulsory. $(5 \times 2 \times 5 = 50)$

Q. 3. (a) Trace the curve $y^2(2a - x) = x^3$.

Ans: The equation of the curve is

$$y^2(2a - x) = x^3 \quad \dots(1)$$

1. **Symmetry:** Since (1) contains only even powers of y , the curve is symmetrical about x axis.

2. **Origin:** The tangents at the origin are given by $y^2 = 0$, i.e., $y = 0$. Since the two tangents are real and coincident, origin is a cusp.

3. **Asymptotes:** Equating to zero, the coefficient of y^2 , the highest degree term in y , the asymptote parallel to y axis is $x - 2a = 0$ i.e., $x = 2a$. There is no other asymptote of the curve.

4. **Points of intersection:** The curve meets x axis and y axis at the origin only.

5. **Region:** From (1), $y = x\sqrt{\frac{x}{2a-x}}$

When $x < 0$, y is imaginary

\Rightarrow No portion of the curve lies to the left of the line $x = 0$ i.e. y axis.

When $0 < x < 2a$, y is real.

When $x > 2a$, y is imaginary

\Rightarrow No portion of curve lies to the right of the line $x = 2a$.

6. **Special points:** From (1),

$$y = \frac{y^{3/2}}{\sqrt{2a-x}} \quad \dots(2)$$

$$\therefore \frac{dy}{dx} = \frac{\sqrt{x}(3a-x)}{(2a-x)^{3/2}}$$

$$\therefore \frac{dy}{dx} = 0$$

$$\text{When } \sqrt{x}(3a-x) = 0$$

$$\Rightarrow x = 0, x = 3a$$

Here $x = 3a$ is not possible, because when $x = 3a$, from (2), y is imaginary.

When $x = 0$, $y = 0$.

\therefore Tangent at $(0, 0)$ i.e. at origin is parallel to x axis i.e. the tangent at origin is x axis.

Again $\frac{dy}{dx} \rightarrow \infty$ when $x \rightarrow 2a$. From (2), $x \rightarrow$

$2a$, $y \rightarrow \infty$. Thus, $x = 2a$ is an asymptote.

When $0 < x < 2a$, $\frac{dy}{dx}$ is positive.

\therefore For positive values of y , y is an increasing function of x , i.e. the curve rises for values of x between 0 and $2a$.

(b) If $Z = f(x + ct) + \phi(x - ct)$ show that

$$\frac{\partial^2 z}{\partial t^2} = c^2 \frac{\partial^2 z}{\partial x^2}.$$

$$\text{Ans. } z = f(x + ct) + \phi(x - ct)$$

$$\Rightarrow \frac{\partial z}{\partial x} = \frac{\partial}{\partial x}[f(x + ct)] + \frac{\partial}{\partial x}[\phi(x - ct)]$$

$$\Rightarrow \frac{\partial z}{\partial x} = f'(x + ct) + \phi'(x - ct)$$

$$\Rightarrow \frac{\partial^2 z}{\partial x^2} = \frac{\partial}{\partial x}[f'(x + ct)] + \frac{\partial}{\partial x}[\phi'(x - ct)]$$

$$\Rightarrow \frac{\partial^2 z}{\partial x^2} = f''(x + ct) + \phi''(x - ct)$$

$$\Rightarrow c^2 \frac{\partial^2 z}{\partial x^2} = c^2 f''(x + ct) + c^2 \phi''(x - ct) \quad \dots(1)$$

$$\text{Also, } \frac{\partial z}{\partial t} = \frac{\partial}{\partial t}[f(x + ct)] + \frac{\partial}{\partial t}[\phi(x - ct)]$$

$$\Rightarrow \frac{\partial z}{\partial t} = f'(x + ct) + \frac{\partial}{\partial t}(ct) + \phi'(x - ct) \cdot \frac{\partial}{\partial t}(-ct)$$

$$\Rightarrow \frac{\partial z}{\partial t} = cf'(x + ct) - c\phi'(x - ct)$$

$$\Rightarrow \frac{\partial^2 z}{\partial t^2} = \frac{\partial}{\partial t}[cf'(x + ct)] - \frac{\partial}{\partial t}[c\phi'(x - ct)]$$

$$\Rightarrow \frac{\partial^2 z}{\partial t^2} = cf''(x + ct) \frac{\partial}{\partial t}(ct) - c\phi''(x - ct) \frac{\partial}{\partial t}(-ct)$$

$$\Rightarrow \frac{\partial^2 z}{\partial t^2} = c^2 f''(x + ct) + c^2 \phi''(x - ct) \quad \dots(2)$$

From (1) and (2),

$$\frac{\partial^2 z}{\partial t^2} = c^2 \frac{\partial^2 z}{\partial x^2}$$

Hence proved.

(c) Expand $e^{ax} \sin by$ in the powers of x and y as far as terms of third degree.

Ans. Given function $f = e^{ax} \sin by$

$$\frac{\partial f}{\partial x} = ae^{ax} \sin by$$

$$\frac{\partial f}{\partial y} = be^{ax} \cos by$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right)$$

$$\Rightarrow \frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} (ae^{ax} \sin by)$$

$$\Rightarrow \frac{\partial^2 f}{\partial x^2} = a^2 e^{ax} \sin by$$

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right) = \frac{\partial}{\partial x} (be^{ax} \cos by)$$

$$\Rightarrow \frac{\partial^2 f}{\partial x \partial y} = abe^{ax} \cos by \quad \frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial y} \right)$$

$$\Rightarrow \frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} (be^{ax} \cos by)$$

$$\Rightarrow \frac{\partial^2 f}{\partial y^2} = -b^2 e^{ax} \sin by$$

$$\frac{\partial^3 f}{\partial x^3} = \frac{\partial}{\partial x} \left(\frac{\partial^2 f}{\partial x^2} \right)$$

$$\Rightarrow \frac{\partial^3 f}{\partial x^3} = \frac{\partial}{\partial x} (a^2 e^{ax} \sin by)$$

$$\Rightarrow \frac{\partial^3 f}{\partial x^3} = a^3 e^{ax} \sin by \quad \frac{\partial^3 f}{\partial x^2 \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial^2 f}{\partial x \partial y} \right)$$

$$\Rightarrow \frac{\partial^3 f}{\partial x^2 \partial y} = \frac{\partial}{\partial x} (abe^{ax} \cos by)$$

$$\Rightarrow \frac{\partial^3 f}{\partial x^2 \partial y} = a^2 be^{ax} \cos by$$

$$\frac{\partial^3 f}{\partial x \partial y^2} = \frac{\partial}{\partial x} \left(\frac{\partial^2 f}{\partial y^2} \right)$$

$$\Rightarrow \frac{\partial^3 f}{\partial x \partial y^2} = \frac{\partial}{\partial x} (-b^2 e^{ax} \sin by)$$

$$\Rightarrow \frac{\partial^3 f}{\partial x \partial y^2} = -ab^2 e^{ax} \sin by \quad \frac{\partial^3 f}{\partial y^3} = \frac{\partial}{\partial y} \left(\frac{\partial^2 f}{\partial y^2} \right)$$

$$\Rightarrow \frac{\partial^3 f}{\partial y^3} = \frac{\partial}{\partial y} (-b^2 e^{ax} \sin by)$$

$$\Rightarrow \frac{\partial^3 f}{\partial y^3} = -b^3 e^{ax} \cos by$$

Now, we make the following table:

Function	Expression	Value at (0, 0)
f	$e^{ax} \sin by$	0
$\frac{\partial f}{\partial x}$	$ae^{ax} \sin by$	0
$\frac{\partial f}{\partial y}$	$be^{ax} \cos by$	b
$\frac{\partial^2 f}{\partial x^2}$	$a^2 e^{ax} \sin by$	0
$\frac{\partial^2 f}{\partial x \partial y}$	$abe^{ax} \cos by$	ab
$\frac{\partial^2 f}{\partial y^2}$	$-b^2 e^{ax} \sin by$	0
$\frac{\partial^3 f}{\partial x^3}$	$a^3 e^{ax} \sin by$	0
$\frac{\partial^3 f}{\partial x^2 \partial y}$	$a^2 be^{ax} \cos by$	$a^2 b$
$\frac{\partial^3 f}{\partial x \partial y^2}$	$-ab^2 e^{ax} \sin by$	0
$\frac{\partial^3 f}{\partial y^3}$	$-b^3 e^{ax} \cos by$	$-b^3$

By Taylor's theorem, expansion of $f(x, y)$ about the point (h, k)

$$\begin{aligned}
 f(x, y) &= f(h, k) + \frac{1}{1!} \left[(x-h) \frac{\partial f}{\partial x} + (y-k) \frac{\partial f}{\partial y} \right] \\
 &+ \frac{1}{2!} \left[(x-h)^2 \frac{\partial^2 f}{\partial x^2} + 2(x-h)(y-k) \frac{\partial^2 f}{\partial x \partial y} + (y-k)^2 \frac{\partial^2 f}{\partial y^2} \right] \\
 &+ \frac{1}{3!} \left[(x-h)^3 \frac{\partial^3 f}{\partial x^3} + 3(x-h)^2(y-k) \frac{\partial^3 f}{\partial x^2 \partial y} \right. \\
 &\left. + 3(x-h)(y-k)^2 \frac{\partial^3 f}{\partial x \partial y^2} + (y-k)^3 \frac{\partial^3 f}{\partial y^3} \right] + \dots
 \end{aligned}$$

Here, $h = 0, k = 0$

$$\begin{aligned}
 \Rightarrow f(x, y) &= f(0, 0) + \frac{1}{1!} \left[x \frac{\partial f}{\partial x} + y \frac{\partial f}{\partial y} \right] + \frac{1}{2!} \left[x^2 \frac{\partial^2 f}{\partial x^2} \right. \\
 &+ 2xy \frac{\partial^2 f}{\partial x \partial y} + y^2 \frac{\partial^2 f}{\partial y^2} \left. \right] + \frac{1}{3!} \left[x^3 \frac{\partial^3 f}{\partial x^3} + 3x^2y \frac{\partial^3 f}{\partial x^2 \partial y} \right. \\
 &\left. + 3xy^2 \frac{\partial^3 f}{\partial x \partial y^2} + y^3 \frac{\partial^3 f}{\partial y^3} \right] + \dots
 \end{aligned}$$

$$\Rightarrow f(x, y) = \frac{1}{1!} [by] + \frac{1}{2!} [ab(2xy)] + \frac{1}{3!} [3x^2y(a^2b) - b_3y_3] + \dots$$

$$\Rightarrow f(x, y) = by + abxy + \frac{1}{2} a^2 bx^2 y - \frac{b^3}{6} y^3 + \dots$$

Q. 4. (a) A rectangular box, open at the top, is to have a volume of 32 cubic feet. Determine the dimensions of the box requiring least material for its construction.

Ans: Let x, y and z be the length, breadth and height of the box respectively.

\therefore volume of the box is 32 cubic feet

$$\Rightarrow xyz = 32 \quad (1)$$

[\therefore volume = length \times breadth \times height]

Let s be the surface area of the box.

$\Rightarrow s$ = Total surface area of closed box – area of the top.

$$= 2(xy + yz + zx) - xy$$

$$[\therefore \text{surface area} = 2(lb + bh + hl)]$$

$$= xy + 2yz + 2zx$$

Here s is to be minimized

$$\therefore \text{Let } f = xy + 2yz + 2zx$$

and $\phi = xyz - 32$

[From (1)]

Lagrange's equations are,

$$\frac{\partial f}{\partial x} + \lambda \frac{\partial \phi}{\partial x} = 0$$

$$\frac{\partial f}{\partial y} + \lambda \frac{\partial \phi}{\partial y} = 0$$

$$\frac{\partial f}{\partial z} + \lambda \frac{\partial \phi}{\partial z} = 0$$

$$\Rightarrow y + 2z + \lambda yz = 0 \quad \dots(2)$$

$$x + 2z + \lambda xz = 0 \quad \dots(3)$$

$$2y + 2x + \lambda xy = 0 \quad \dots(4)$$

$$\text{From (2), } \lambda = -\frac{y+2z}{yz} \quad \dots(5)$$

$$\text{From (3), } \lambda = -\frac{x+2z}{xz} \quad \dots(6)$$

$$\text{From (4), } \lambda = -\frac{2x+2y}{xy} \quad \dots(7)$$

Equating λ from (5) and (6), we get,

$$-\frac{y+2z}{yz} = -\frac{x+2z}{xz}$$

$$\Rightarrow x(y+2z) = y(x+2z)$$

$$\Rightarrow xy + 2xz = xy + 2yz$$

$$\Rightarrow 2xz = 2yz$$

$$\Rightarrow x = y$$

Again, equating λ from (5) and (7), we get,

$$-\frac{y+2z}{yz} = -\frac{2x+2y}{xy}$$

$$\Rightarrow x(y+2z) = z(2x+2y)$$

$$\Rightarrow xy + 2xz = 2xz + 2yz$$

$$\Rightarrow xy = 2yz$$

$$\Rightarrow z = \frac{x}{2}$$

Putting $y = x$ and $z = \frac{x}{2}$ in (1), we get,

$$(x)(x)\left(\frac{x}{2}\right) = 32$$

$$\Rightarrow x^3 = 64$$

$$\Rightarrow x = 4$$

$$\Rightarrow y = 4 \quad [\because y = x]$$

$$\Rightarrow z = 2 \quad [\because z = \frac{x}{2}]$$

Thus, length, breadth and height of the box are 4 feet, 4 feet and 2 feet respectively, requiring least material for the construction of the box.

(b) If $u_1 = \frac{x_2 x_3}{x_1}$, $u_2 = \frac{x_3 x_1}{x_2}$ and $u_3 = \frac{x_1 x_2}{x_3}$ find

the value of $\frac{\partial(u_1, u_2, u_3)}{\partial(x_1, x_2, x_3)}$.

Ans: We have

$$\frac{\partial(u_1, u_2, u_3)}{\partial(x_1, x_2, x_3)} = \begin{vmatrix} \frac{\partial u_1}{\partial x_1} & \frac{\partial u_1}{\partial x_2} & \frac{\partial u_1}{\partial x_3} \\ \frac{\partial u_2}{\partial x_1} & \frac{\partial u_2}{\partial x_2} & \frac{\partial u_2}{\partial x_3} \\ \frac{\partial u_3}{\partial x_1} & \frac{\partial u_3}{\partial x_2} & \frac{\partial u_3}{\partial x_3} \end{vmatrix}$$

$$= \begin{vmatrix} \frac{x_2 x_3}{x_1^2} & \frac{x_3}{x_1} & \frac{x_2}{x_1} \\ \frac{x_3}{x_2} & -\frac{x_1 x_3}{x_2^2} & \frac{x_1}{x_2} \\ \frac{x_2}{x_3} & \frac{x_1}{x_3} & -\frac{x_1 x_2}{x_3^2} \end{vmatrix}$$

$$= \frac{1}{x_1^2} \cdot \frac{1}{x_2^2} \cdot \frac{1}{x_3^2} \begin{vmatrix} -x_2 x_3 & x_1 x_3 & x_1 x_2 \\ x_2 x_3 & -x_1 x_3 & x_1 x_2 \\ x_2 x_3 & x_1 x_3 & -x_1 x_2 \end{vmatrix}$$

[Taking $\frac{1}{x_1^2}$ from R_1 , $\frac{1}{x_2^2}$ from R_2 and $\frac{1}{x_3^2}$ from R_3 common]

$$= \frac{(x_2 x_3)(x_1 x_3)(x_1 x_2)}{x_1^2 x_2^2 x_3^2} \begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix}$$

[Taking $x_2 x_3$ from C_1 , $x_1 x_3$ from C_2 and $x_1 x_2$ from C_3 common]

$$= \frac{x_1^2 x_2^2 x_3^2}{x_1^2 x_2^2 x_3^2} \begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix}$$

$$= -1[1 - 1] - 1[-1 - 1] + 1[1 + 1]$$

$$= 4$$

(c) Find the percentage of error in calculating

the area of an ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, when error of + 1% is made in measuring the major and minor axes.

Ans: Eqⁿ of given ellipse is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

\therefore error in major axis and minor axis are 1%.

$$\Rightarrow \frac{\delta a}{a} = \frac{1}{100} \text{ and } \frac{\delta b}{b} = \frac{1}{100}$$

Let A represents the area of ellipse. To find error in A, we shall express A in terms of a and b, because errors in a and b are known to us. We know that

$A = \pi ab$ [\therefore area of ellipse = πxy where x and y are semimajor and semiminor axis]

Taking log of both sides, we get,

$$\log A = \log(\pi ab)$$

$$\Rightarrow \log A = \log \pi + \log a + \log b$$

$$[\because \log(mn) = \log m + \log n]$$

By differentials, we get,

$$\frac{\delta A}{A} = 0 + \frac{\delta a}{a} + \frac{\delta b}{b} \quad \left[\begin{array}{l} \therefore \pi \text{ is constant} \\ \therefore \text{its differential is zero} \end{array} \right]$$

$$\Rightarrow \frac{\delta A}{A} = \frac{\delta a}{a} + \frac{\delta b}{b}$$

$$\Rightarrow \frac{\delta A}{A} = \frac{1}{100} + \frac{1}{100} \quad \left[\because \frac{\delta a}{a} = \frac{1}{100}, \frac{\delta b}{b} = \frac{1}{100} \right]$$

$$\Rightarrow \frac{\delta A}{A} = \frac{2}{100}$$

Thus, error in area of ellipse is 2%.

Q. 5. (a) Test for consistency and solve the following system of equations

$$2x - y + 3z = 8$$

$$-x + 2y + z = 4$$

$$3x + y - 4z = 0$$

Ans: Given system of equations is

$$2x - y + 3z = 8$$

$$\begin{aligned} -x + 2y + z &= 4 \\ 3x + y - 4z &= 0 \end{aligned}$$

$$\text{Augmented matrix } [A : B] = \begin{bmatrix} 2 & -1 & 3 & : & 8 \\ -1 & 2 & 1 & : & 4 \\ 3 & 1 & -4 & : & 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} -1 & 2 & 1 & : & 4 \\ 2 & -1 & 3 & : & 8 \\ 3 & 1 & -4 & : & 0 \end{bmatrix} R_1 \leftrightarrow R_2$$

$$\sim \begin{bmatrix} -1 & 2 & 1 & : & 4 \\ 0 & 3 & 5 & : & 16 \\ 0 & 7 & -1 & : & 12 \end{bmatrix} \begin{aligned} R_2 &\rightarrow R_2 + 2R_1 \\ R_3 &\rightarrow R_3 + 3R_1 \end{aligned}$$

$$\sim \begin{bmatrix} -1 & 2 & 1 & : & 4 \\ 0 & -1 & 11 & : & 20 \\ 0 & 7 & -1 & : & 12 \end{bmatrix} R_2 \rightarrow 2R_2 - R_3$$

$$\sim \begin{bmatrix} -1 & 2 & 1 & : & 4 \\ 0 & -1 & 11 & : & 20 \\ 0 & 0 & 76 & : & 152 \end{bmatrix} R_3 \rightarrow R_3 + 7R_2$$

which is in echelon form.

Here, rank of $A = 3$

Rank of $[A : B] = 3$

No of unknowns = 3

\Rightarrow Rank of $A =$ Rank of $[A : B] =$ No. of unknowns

\Rightarrow System is consistent with unique solution.

Rewriting the system,

$$-x + 2y + z = 4 \quad \dots(1)$$

$$-y + 11z = 20 \quad \dots(2)$$

$$76z = 152 \quad \dots(3)$$

$$\text{From (3)} \quad z = \frac{152}{76} = 2$$

Put $z = 2$ in (2), we get

$$-y + 11(2) = 20$$

$$\Rightarrow -y = -2 \Rightarrow y = 2$$

Putting $y = 2, z = 2$ in (1), we get,

$$-x + 2(2) + 2 = 4$$

$$\Rightarrow -x + 6 = 4 \Rightarrow x = 2$$

Thus, $x = 2, y = 2, z = 2$ Ans.

(b) Reduce the following matrix to normal form and hence find its rank:

$$\begin{bmatrix} 5 & 3 & 14 & 4 \\ 0 & 1 & 2 & 1 \\ 1 & -1 & 2 & 0 \end{bmatrix}$$

$$\text{Ans: Given matrix} = \begin{bmatrix} 5 & 3 & 14 & 4 \\ 0 & 1 & 2 & 1 \\ 1 & -1 & 2 & 0 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & -1 & 2 & 0 \\ 0 & 1 & 2 & 1 \\ 5 & 3 & 14 & 4 \end{bmatrix} R_1 \leftrightarrow R_3$$

$$\sim \begin{bmatrix} 1 & -1 & 2 & 0 \\ 0 & 1 & 2 & 1 \\ 0 & 8 & 4 & 4 \end{bmatrix} R_3 \rightarrow R_3 - 5R_1$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 \\ 0 & 8 & 4 & 4 \end{bmatrix} \begin{aligned} c_2 &\rightarrow c_2 + c_1 \\ c_3 &\rightarrow c_3 - 2c_1 \end{aligned}$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & -12 & -4 \end{bmatrix} R_3 \rightarrow R_3 - 8R_2$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -12 & -4 \end{bmatrix} \begin{aligned} c_3 &\rightarrow c_3 - 2c_2 \\ c_4 &\rightarrow c_4 - c_2 \end{aligned}$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -4 \end{bmatrix} c_3 \rightarrow \frac{c_3}{-12}$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} c_4 \rightarrow c_4 + 4c_3$$

$$\sim [I_3 \ 0]$$

which is in normal form.

\therefore Rank = order of identity matrix in normal form = 3.

(c) Show that the matrix

$$\begin{bmatrix} a+ic & -b+id \\ b+id & a-ic \end{bmatrix}$$

is unitary if and only if $a^2 + b^2 + c^2 + d^2 = 1$.

Ans: Given matrix $A = \begin{bmatrix} a+ic & -b+id \\ b+id & a-ic \end{bmatrix}$. Now $\bar{A} = \begin{bmatrix} a-ic & -b-id \\ b-id & a+ic \end{bmatrix}$ [Replacing i by $-i$]

$$\Rightarrow A^0 = (\bar{A})^T \Rightarrow A^0 = \begin{bmatrix} a-ic & b-id \\ -b-id & a+ic \end{bmatrix}$$

Now $AA^0 = \begin{bmatrix} a+ic & -b+id \\ b+id & a-ic \end{bmatrix} \begin{bmatrix} a-ic & b-id \\ -b-id & a+ic \end{bmatrix}$

$$\Rightarrow AA^0 = \begin{bmatrix} (a+ic)(a-ic) + (-b+id)(-b-id) & (a+ic)(b-id) + (-b+id)(a+ic) \\ (b+id)(a-ic) + (a-ic)(-b-id) & (b+id)(b-id)(a-ic)(a+ic) \end{bmatrix}$$

$$= \begin{bmatrix} a^2 - i^2 c^2 + b^2 - i^2 d^2 & (a+ic)(b-id) - (b-id)(a+ic) \\ (b+id)(a-ic) - (a-ic)(b+id) & b^2 - i^2 d^2 + a^2 - i^2 c^2 \end{bmatrix} = \begin{bmatrix} a^2 + c^2 + b^2 + d^2 & 0 \\ 0 & b^2 + d^2 + a^2 + c^2 \end{bmatrix}$$

Now above matrix is identity matrix $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ if and only if $a^2 + b^2 + c^2 + d^2 = 1$

i.e. $AA^0 = I$ if and only if $a^2 + b^2 + c^2 + d^2 = 1$ i.e. A is unitary if and only if $a^2 + b^2 + c^2 + d^2 = 1$

Q. 6. (a) Prove that $\int_0^1 \frac{dx}{\sqrt{1+x^4}} = \frac{1}{4\sqrt{2}} \beta\left(\frac{1}{4}, \frac{1}{2}\right)$

Ans: Given integral $I = \int_0^1 \frac{dx}{\sqrt{1+x^4}}$

Put $x^4 = \tan^2 \theta$

$$\Rightarrow x = \sqrt{\tan \theta} \Rightarrow dx = \frac{1}{2\sqrt{\tan \theta}} \sec^2 \theta d\theta$$

at $x = 0$, $\tan \theta = 0 \Rightarrow \theta = 0$, at $x = 1$, $\tan \theta = 1 \Rightarrow \theta = \frac{\pi}{4}$

Thus, $I = \int_0^{\frac{\pi}{4}} \frac{1}{\sqrt{1+\tan^2 \theta}} \cdot \frac{1}{2\sqrt{\tan \theta}} \sec^2 \theta d\theta = \frac{1}{2} \int_0^{\frac{\pi}{4}} \frac{1}{\sec \theta} \cdot \frac{1}{\sqrt{\tan \theta}} \sec^2 \theta d\theta$ [$\because 1 + \tan^2 \theta = \sec^2 \theta$]

$$= \frac{1}{2} \int_0^{\frac{\pi}{4}} \sqrt{\cot \theta} \sec \theta d\theta = \frac{1}{2} \int_0^{\frac{\pi}{4}} \frac{\sqrt{\cos \theta}}{\sqrt{\sin \theta} \cos \theta} d\theta = \frac{1}{2} \int_0^{\frac{\pi}{4}} \frac{d\theta}{\sqrt{\sin \theta \cos \theta}} = \frac{\sqrt{2}}{2} \int_0^{\frac{\pi}{4}} \frac{d\theta}{\sqrt{2 \sin \theta \cos \theta}}$$

$$= \frac{\sqrt{2}}{2} \int_0^{\frac{\pi}{4}} \frac{d\theta}{\sqrt{\sin 2\theta}} = \frac{\sqrt{2}}{2} \int_0^{\frac{\pi}{4}} \frac{1}{2} \frac{d\theta}{\sqrt{\sin \phi}}$$

$$\begin{cases} \text{Put } 2\theta = \phi \\ \Rightarrow d\theta = \frac{d\phi}{2} \\ \text{at } \theta = 0, \phi = 0 \\ \text{at } \theta = \frac{\pi}{4}, \phi = \frac{\pi}{2} \end{cases}$$

$$= \frac{\sqrt{2}}{4} \int_0^{\pi/2} \sin^{-1/2} \phi d\phi$$

$$= \frac{\sqrt{2}}{4} \int_0^{\pi} \sin^{-\frac{1}{2}} \phi \cos^0 \phi d\phi = \frac{\sqrt{2}}{4} \left[\frac{-\frac{1}{2}+1}{2} \left| \frac{0+1}{2} \right| \right]$$

$$\left[\because \int_0^{\pi} \sin^p \theta \cos^q \theta d\theta = \frac{\frac{p+1}{2} \left| \frac{q+1}{2} \right|}{2 \left| \frac{p+q+2}{2} \right|} \right]$$

$$= \frac{\sqrt{2}}{8} \left[\frac{1}{4} \left| \frac{1}{2} \right| \right] = \frac{\sqrt{2}}{8} \left[\frac{1}{4} \cdot \frac{1}{2} \right]$$

$$= \frac{\sqrt{2}}{8} \beta\left(\frac{1}{4}, \frac{1}{2}\right) \left[\because \beta(m, n) = \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n)} \right]$$

$$= \frac{1}{4\sqrt{2}} \beta\left(\frac{1}{4}, \frac{1}{2}\right)$$

$$\Rightarrow \int_0^1 \frac{dx}{\sqrt{1+x^4}} = \frac{1}{4\sqrt{2}} \beta\left(\frac{1}{4}, \frac{1}{2}\right)$$

Hence proved.

(b) Evaluate $\iiint_V x^{l-1} y^{m-1} z^{n-1} dx dy dz$, where $x > 0, y > 0$ under the condition

$$\left(\frac{x}{a}\right)^p + \left(\frac{y}{b}\right)^q + \left(\frac{z}{c}\right)^r \leq 1.$$

Ans: Given integral $I = \iiint_V x^{l-1} y^{m-1} z^{n-1} dx dy dz$

where v is the region $x > 0, y > 0, z > 0$

$$\text{and } \left(\frac{x}{a}\right)^p + \left(\frac{y}{b}\right)^q + \left(\frac{z}{c}\right)^r \leq 1$$

$$\text{Put } \left(\frac{x}{a}\right)^p = X, \left(\frac{y}{b}\right)^q = Y, \left(\frac{z}{c}\right)^r = Z$$

$$\Rightarrow \frac{x}{a} = X^{\frac{1}{p}}, \frac{y}{b} = Y^{\frac{1}{q}}, \frac{z}{c} = Z^{\frac{1}{r}}$$

$$\Rightarrow x = a X^{\frac{1}{p}}, y = b Y^{\frac{1}{q}}, z = c Z^{\frac{1}{r}}$$

$$\Rightarrow dx = \frac{a}{p} X^{\frac{1}{p}-1} dX, dy = \frac{b}{q} Y^{\frac{1}{q}-1} dY, dz = \frac{c}{r} Z^{\frac{1}{r}-1} dZ$$

Putting above values,

$$I = \iiint_V \left(a X^{\frac{1}{p}} \right)^{l-1} \left(b Y^{\frac{1}{q}} \right)^{m-1} \left(c Z^{\frac{1}{r}} \right)^{n-1} \frac{a}{p} \frac{b}{q} \frac{c}{r} dX dY dZ$$

$$X^{\frac{l-1}{p}} Y^{\frac{m-1}{q}} Z^{\frac{n-1}{r}} dX dY dZ$$

where v is the region $Z > 0, Y > 0, X > 0$

and $X + Y + Z \leq 1$.

$$\Rightarrow I = \frac{1}{pqr} \iiint_V X^{\frac{l-1}{p}} Y^{\frac{m-1}{q}} Z^{\frac{n-1}{r}} dX dY dZ$$

$$X^{\frac{l-1}{p}} Y^{\frac{m-1}{q}} Z^{\frac{n-1}{r}} dX dY dZ$$

$$= \frac{a^l b^m c^n}{pqr} \iiint_V X^{\frac{l-1}{p}} Y^{\frac{m-1}{q}} Z^{\frac{n-1}{r}} dX dY dZ$$

$$= \frac{a^l b^m c^n}{pqr} \iiint_V X^{\frac{l-1}{p}} Y^{\frac{m-1}{q}} Z^{\frac{n-1}{r}} dX dY dZ$$


$$= \frac{a^l b^m c^n}{pqr} \frac{\left| \frac{l}{p} \right| \left| \frac{m}{q} \right| \left| \frac{n}{r} \right|}{\left| \frac{l}{p} + \frac{m}{q} + \frac{n}{r} + 1 \right|} \text{ Ans.}$$

[\therefore By Dirichlet's theorem,

$$\iiint_V x^{l-1} y^{m-1} z^{n-1} dx dy dz = \frac{\Gamma(l) \Gamma(m) \Gamma(n)}{\Gamma(l+m+n+1)}]$$

(c) Find the area of one loop of the lemniscates $r^2 = a^2 \cos 2\theta$.

Ans: The given lemniscate is shown in the following figure.



Required area = $\iint_R r dr d\theta$ where R is the region bounded by one loop of lemniscate $r^2 = a^2 \cos 2\theta$.

Limits of θ are 0 to $a \sqrt{\cos 2\theta}$

Limits of r are $-\frac{\pi}{4}$ to $\frac{\pi}{4}$.

$$\begin{aligned} \text{Thus, required area} &= \int_{-\pi/4}^{\pi/4} \int_0^{a\sqrt{\cos 2\theta}} r dr d\theta \\ &= \int_{-\pi/4}^{\pi/4} \left[\frac{r^2}{2} \right]_{r=0}^{a\sqrt{\cos 2\theta}} d\theta = \frac{1}{2} \int_{-\pi/4}^{\pi/4} a^2 \cos^2 \theta d\theta \\ &= \frac{a^2}{2} \left[\frac{\sin 2\theta}{2} \right]_{-\pi/4}^{\pi/4} \left[\because \int \cos ax dx = \frac{\sin ax}{a} \right] \\ &= \frac{a^2}{4} \left[\sin \frac{\pi}{2} - \sin \left(-\frac{\pi}{2} \right) \right] \\ &= \frac{a^2}{4} [1 + 1] = \frac{a^2}{2} \text{ Ans.} \end{aligned}$$

Q. 7. (a) Find the directional derivative of $\phi(x, y, z) = xy^2 + yz^3$ at the point $(2, -1, 1)$ in the direction of the normal to the surface $x \log z - y^2 + 4 = 0$ at $(2, -1, 1)$

Ans: Given surface $\phi = xy^2 + yz^3$

$$\begin{aligned} \text{Now } \nabla \phi &= \hat{i} \frac{\partial \phi}{\partial x} + \hat{j} \frac{\partial \phi}{\partial y} + \hat{k} \frac{\partial \phi}{\partial z} \\ \Rightarrow \nabla \phi &= \hat{i}(y^2) + \hat{j}(2xy + z^3) + \hat{k}(3yz^2) \\ \Rightarrow (\nabla \phi)_{(2, -1, 1)} &= \hat{i} + \hat{j}(-4 + 1) + \hat{k}(-3) \\ \Rightarrow (\nabla \phi)_{(2, -1, 1)} &= \hat{i} - 3\hat{j} - 3\hat{k} \\ \therefore \text{ We want to find normal to the surface } &x \log z - y^2 + 4 = 0 \\ \therefore \text{ Let } g &= x \log z - y^2 + 4 \end{aligned}$$

$$\Rightarrow \nabla g = \frac{\partial g}{\partial x} \hat{i} + \frac{\partial g}{\partial y} \hat{j} + \frac{\partial g}{\partial z} \hat{k}$$

$$\Rightarrow \nabla g = \log z \hat{i} - 2y \hat{j} + \frac{x}{z} \hat{k}$$

$$\Rightarrow (\nabla g)_{(2, -1, 1)} = \log 1 \hat{i} + 2\hat{j} + \frac{2}{1} \hat{k}$$

$$\Rightarrow \vec{n} = 2\hat{j} + 2\hat{k}$$

[$\because \log 1 = 0$ and $\vec{n} = \nabla g$ represent normal to surface]

$$\Rightarrow \hat{n} = \frac{\vec{n}}{|\vec{n}|}$$

$$\Rightarrow \hat{n} = \frac{1}{\sqrt{2^2 + 2^2}} (2\hat{j} + 2\hat{k})$$

$$\left[\because \vec{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k} \right. \\ \left. \Rightarrow |\vec{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2} \right] = \frac{1}{2\sqrt{2}} (2\hat{j} + 2\hat{k})$$

$$\Rightarrow \hat{n} = \frac{1}{\sqrt{2}} (\hat{j} + \hat{k})$$

Now, Desired directional derivative = $\nabla \phi \cdot \hat{n}$

$$= (1 - 3\hat{j} - 3\hat{k}) \cdot \frac{1}{\sqrt{2}} (\hat{j} + \hat{k}) = \frac{1}{\sqrt{2}} (-3 - 3)$$

$$\left[\begin{aligned} \because \vec{a} &= a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k} \\ \vec{b} &= b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k} \\ \Rightarrow \vec{a} \cdot \vec{b} &= a_1 b_1 + a_2 b_2 + a_3 b_3 \end{aligned} \right]$$

$$= \frac{-6}{\sqrt{2}} = \frac{-6}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} = -3\sqrt{2} \text{ Ans.}$$

(b) If all second order derivatives of ϕ and \vec{v} are continuous, then show that

$$(i) \text{ curl } (\text{grad } \phi) = \vec{v}$$

$$(ii) \text{ div } (\text{curl } \vec{v}) = 0$$

Ans: (i) To show that $\text{curl } (\text{grad } \phi) = 0$, we shall first find $\text{grad } \phi$.

Now, $\text{grad } \phi = \nabla \phi$

$$\Rightarrow \text{grad } \phi = \hat{i} \frac{\partial \phi}{\partial x} + \hat{j} \frac{\partial \phi}{\partial y} + \hat{k} \frac{\partial \phi}{\partial z}$$

Now $\text{curl } \vec{v} = \nabla \times \vec{v}$

$$\Rightarrow = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ V_1 & V_2 & V_3 \end{vmatrix}$$

$$\Rightarrow \text{curl}(\text{grad } \phi) = \nabla \times (\nabla \phi)$$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{\partial \phi}{\partial x} & \frac{\partial \phi}{\partial y} & \frac{\partial \phi}{\partial z} \end{vmatrix}$$

$$= \hat{i} \left[\frac{\partial}{\partial y} \left(\frac{\partial \phi}{\partial z} \right) - \frac{\partial}{\partial z} \left(\frac{\partial \phi}{\partial y} \right) \right] + \text{two similar terms}$$

$$= \hat{i} \left[\frac{\partial^2 \phi}{\partial y \partial z} - \frac{\partial^2 \phi}{\partial y \partial z} \right] + \text{two similar terms} = 0$$

$$\Rightarrow \text{curl}(\text{grad } \phi) = 0$$

Hence proved.

(ii) To show that $\text{div}(\text{curl } \vec{v}) = 0$, we shall first find $\text{curl } \vec{v}$.

$$\text{Let } \vec{v} = v_1 \hat{i} + v_2 \hat{j} + v_3 \hat{k}$$

$$\text{curl } \vec{v} = \nabla \times \vec{v} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ V_1 & V_2 & V_3 \end{vmatrix}$$

$$\Rightarrow \text{curl } \vec{v} = \hat{i} \left(\frac{\partial v_3}{\partial y} - \frac{\partial v_2}{\partial z} \right) + \hat{j} \left(\frac{\partial v_1}{\partial z} - \frac{\partial v_3}{\partial x} \right) + \hat{k} \left(\frac{\partial v_2}{\partial x} - \frac{\partial v_1}{\partial y} \right)$$

$$\text{Now, } \text{div } \vec{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z}$$

$$\Rightarrow \text{div}(\text{curl } \vec{v}) = \frac{\partial}{\partial x} \left(\frac{\partial v_3}{\partial y} - \frac{\partial v_2}{\partial z} \right) + \frac{\partial}{\partial y} \left(\frac{\partial v_1}{\partial z} - \frac{\partial v_3}{\partial x} \right) + \frac{\partial}{\partial z} \left(\frac{\partial v_2}{\partial x} - \frac{\partial v_1}{\partial y} \right)$$

$$= \frac{\partial^2 V_3}{\partial x \partial y} - \frac{\partial^2 V_2}{\partial x \partial z} + \frac{\partial^2 V_1}{\partial y \partial z} - \frac{\partial^2 V_3}{\partial x \partial y}$$

$$= \frac{\partial^2 V_2}{\partial x \partial z} - \frac{\partial^2 V_1}{\partial y \partial z} = 0$$

$$\Rightarrow \text{div}(\text{curl } \vec{v}) = 0 \text{ Hence Proved.}$$

(c) Find the work done by the force

$$\vec{f} = (2y + 3)\hat{i} + xz\hat{j} + (yz - x)\hat{k}$$

when it moves a particle from the point (0, 0, 0) to the point (2, 1, 1) along the curve $x = 2t^2$, $y = t$ and $z = t^3$,

$$\text{Ans. Required work done} = \int_c \vec{f} \cdot d\vec{r}$$

$$= \int_c [(dy + 3)dx + xzdy + (yz - x)dz] \quad \dots(1)$$

$$\left[\begin{aligned} \because \vec{F} &= F_1 \hat{i} + F_2 \hat{j} + F_3 \hat{k}, d\vec{r} = dx \hat{i} + dy \hat{j} + dz \hat{k} \\ \Rightarrow \vec{F} \cdot d\vec{r} &= F_1 dx + F_2 dy + F_3 dz \end{aligned} \right]$$

Along c,

$$x = 2t^2, y = t, z = t^3$$

$$\Rightarrow dx = 4t dt, dy = dt, dz = 3t^2 dt$$

t varies from 0 to 1.

Putting these values in (1), we get,

Required work done

$$= \int_{t=0}^1 (2t + 3)4t dt + \int_{t=0}^1 (2t^2)(t^3) dt + \int_{t=0}^1 (t \cdot t^3 - 2t^2)3t^2 dt$$

$$= \int_{t=0}^1 (8t^2 + 12t + 2t^5 + 3t^6 - 6t^4) dt$$

$$= \left[\frac{8t^3}{3} + 12 \frac{t^2}{2} + 2 \frac{t^6}{6} + \frac{3t^7}{7} - \frac{6t^5}{5} \right]_{t=0}^1$$

$$= \frac{8}{3} + \frac{12}{2} + \frac{2}{6} + \frac{3}{7} - \frac{6}{5}$$

$$= \frac{560 + 1260 + 70 + 90 - 252}{210}$$

$$= \frac{1728}{210} = \frac{288}{35} \text{ Ans.}$$