| Enrol | lment | No. |
|-------|-------|-----|
|       |       |     |

#### Seat No.: \_

## **GUJARAT TECHNOLOGICAL UNIVERSITY**

**B.E. Sem-I** Examination January 2010

Subject code: 110008 Date: 11 / 01 /2010

Subject Name: Mathematics – I Time: 11.00 am – 02.00 pm Total Marks: 70

### **Instructions:**

- 1. Attempt all questions.
- 2. Make suitable assumptions wherever necessary.
- 3. Figures to the right indicate full marks.
- Find the value of k so that the function given below is continuous at a given Q1. (i) **(a)** point x=2.

$$f(x) = \begin{cases} \frac{2^{x+2} - 16}{4^x - 16}, & x \neq 2\\ k, & x = 2 \end{cases}$$

- 02 (ii) State Sandwich theorem and using it find  $\lim_{x \to a} g(x)$  if  $3 - x^3 \le g(x) \le 3 \sec x$ for all x.
- If f(x) and g(x) are continuous functions for  $0 \le x \le 1$ , could **(b)** (i) 02 f(x)/g(x) possibly be discontinuous at a point in the interval [0,1]? Give reasons for your answer.
  - If  $f:(a,b) \to \Re$  is differentiable at  $c \in (a,b)$ , then show that 02 (ii)

 $\lim_{h \to 0^+} \frac{f(c+a) - f(c+b)}{2h}$  exists and equals f'(c). Is the converse true?

(c) (i) Using Mean Value Theorem, Prove 
$$0 < \frac{1}{x} \log \left( \frac{e^x - 1}{x} \right) < 1$$
, for  $x > 0$ . 03

(ii) For what values of 
$$a, m$$
 and  $b$  does the function 03  
 $\begin{cases} 3 \\ 2 \end{cases}, x = 0 \end{cases}$ 

$$f(x) = \begin{cases} -x^2 + 3x + a, \ 0 < x < 1 \\ mx + b, \ 1 \le x \le 2 \end{cases}$$

satisfy the hypothesis of the Mean Value Theorem on the interval [0,2].

Q2. (a) (i) Find the area of the region between the *x*-axis and the graph of 
$$f(x) = x^3 - x^2 - 2x, -1 \le x \le 2$$
.

Using Fundamental Theorem of Calculus find  $\frac{dy}{dx}$  if  $y = \int_{1}^{x^2} \cos t \, dt$ . 02 **(ii)** 

- Evaluate the integral  $\int_{0}^{\infty} \frac{dx}{x^2+1}$ . 02 (iii)
- Find the absolute maximum and minimum values of the function on the given 03 **(b)** (i) interval  $f(t) = |t-5|, 4 \le t \le 7$ .
  - (ii) 02 Find the Taylor's series expansion of  $f(x) = x^3 - 2x + 4$ , a = 2.

1

02

## Download all NOTES and PAPERS at StudentSuvidha.com

(iii) The geometric mean of two positive numbers *a* and *b* is the number  $\sqrt{ab}$ . 02 Show that the value of *c* in the conclusion of the Mean Value Theorem for  $f(x) = \frac{1}{x}$  on an interval of positive numbers [a,b] is  $c = \sqrt{ab}$ .

#### OR

(b) (i) Test the convergence or divergence of the following series (ANY TWO) 04

**a.** 
$$\sum_{n=0}^{\infty} \frac{2^n - 1}{3^n}$$
 **b.**  $\sum_{n=1}^{\infty} \frac{n}{e^{-n}}$  **c.**  $\sum_{n=0}^{\infty} n! (x-4)^n$ 

(ii) Using Riemann Sum show that 
$$\int_{a}^{b} x \, dx = \frac{1}{2} (b^2 - a^2)$$
 03

Q3. (a) Suppose that 
$$w = f(x, y)$$
,  $x = g(r, s)$  and  $y = h(r, s)$  then write the chain rule 05  
for  $\frac{\partial w}{\partial r}$  and  $\frac{\partial w}{\partial s}$ . Also evaluate  $\frac{\partial w}{\partial r}$  and  $\frac{\partial w}{\partial s}$  in terms of r and s if  
 $w = x + 2y + z^2$ ,  $x = \frac{r}{s}$ ,  $y = r^2 + \ln s$ ,  $z = 2r$ .  
(b) (i) Show that  $\int_{-\infty}^{\infty} f(x) dx$  may not equal to  $\lim_{b \to \infty} \int_{b}^{b} f(x) dx$ . 02

(ii) If 
$$u = \tan^{-1}\left(\frac{x^2 + y^2}{x - y}\right)$$
 show that  $x\frac{\partial u}{\partial x} + y\frac{\partial u}{\partial y} = \frac{1}{2}\sin 2u$  (03)

(c) Find the length of the curve 
$$y = \int_{0}^{1} \sqrt{\cos 2t} dt$$
 from  $x = 0$  to  $x = \frac{\pi}{4}$ . 04

#### 0

from C

Q3. (a) Let 
$$y = f(x, y, z)$$
 be a function of three independent variables, write the 05  
formal definition of the partial derivative for  $\frac{\partial f}{\partial z}$  at  $(x_0, y_0, z_0)$ . Using this definition find  $\frac{\partial f}{\partial z}$  at  $(1, 2, 3)$  for  $f(x, y, z) = x^2 y z^2$ .  
(b) (i) Show that 03  
 $f(x, y) = \begin{cases} \frac{2xy}{x^2 + y^2}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$  is continuous at every point except at the origin.

(ii) Find  $\frac{dw}{dt}$  if w = xy + z,  $x = \cos t$ ,  $y = \sin t$ , z = t. 02

(c) Find the volume of the solid generated by revolving the region bounded by  $y = \sqrt{x}$  and the lines y = 2 and x = 0 about the line y = 2.

Q4. (a) (i) Evaluate the integral 
$$\int_{0}^{1} \int_{0}^{\sqrt{1-y^2}} \int_{0}^{2} (x^2 + y^2) dx dy$$
 03

(ii) Find the volume of the region that lies under the paraboloid  $z = x^2 + y^2$  and above the triangle enclosed by the lines y = x, x = 0 and x + y = 2 in the xy - y plane.

#### 2

## Download all NOTES and PAPERS at StudentSuvidha.com

| <b>(b)</b> |   | Find the equations for tangent plane and normal line at the point $(1,1,1)$ on the  | 03   |
|------------|---|---|--|
| (c)        |   | surface $x^2 + y^2 + z^2 = 3$ .<br>Find the area of the region that lies inside the cardioid $r = 1 + \cos \vartheta$ and outside the circle $r = 1$ .  | 04   |
|            |   | OR  |  |
| (a)        |   | Evaluate $\int_{0}^{1} \int_{0}^{\sqrt{z}} \int_{0}^{2\pi} (r^2 \cos^2 \vartheta + z^2) r  d\vartheta  dr  dz  .$   | 04   |
| (b)        | (i)   | Integrate $f(x, y) = \frac{\ln(x^2 + y^2)}{\sqrt{x^2 + y^2}}$ over the region $1 \le x^2 + y^2 \le e$ by changing   | 04   |
|            |   | to polar coordinates  |  |
|            | (ii)  | Find the derivative of $f(x, y, z) = x^3 - xy^2 - z$ at $P_0(1, 1, 0)$ in the direction   | 02   |
|            |   | $\rightarrow$ $\rightarrow$ $\rightarrow$ $\rightarrow$ $\rightarrow$   |  |
| (a)        |   | of $V = 2I - 5J + 6K$ .<br>Find the volume of the prism whose base is the triangle in $m_{1}$ plane   | 04   |
| (0)        |   | Find the volume of the prism whose base is the triangle in $xy$ - plane<br>bounded by the x - axis and the line $y = x$ and $x = 1$ and whose top lies in   | 04   |
|            |   | the plane $z = f(x, y) = 3 - x - y$   |  |
| (9)        |   | $\int \int \frac{1}{\sqrt{2}} dx = \int \frac{1}{\sqrt$ | 05   |
| (4)        |   | Integrate $f(x, y, z) = x + \sqrt{y - z^2}$ over the path $C = C_1 \cup C_2$ from   | 00   |
|            |   | (0,0,0) to $(1,1,1)$ with   |  |
|            |   | $C_1: r(t) = t \vec{i} + t^2 \vec{j}, \ 0 \le t \le 1$  |  |
|            |   | $C : v(t) = i + i + tk  0 \le t \le 1$  |  |
| (h)        |   | $C_2 \cdot f(t) = t + j + t k, 0 \le t \le 1$   | 05   |
| (0)        |   | State Green's theorem and also evaluate the integral $\oint (6y + x)dx + (y + 2x)dy$  | 03   |
|            |   | $1 - (2 - 1)^2 + (2 - 2)^2 - 1$   |  |
| (a)        |   | where $C_{x}$ the circle $(x-2)^{2} + (y-3)^{2} = 4$ .  | 04   |
| (C)        |   | Trace the curve $r^2 = a^2 \cos 2\theta$ .  | 04   |
|            |   | OR (  |  |
| <b>(a)</b> | Ń   | Use Green's theorem to evaluate the integral $\oint (y^2 dx + x^2 dy)$ where  | 05   |
|            | •   |   |  |
|            |   | The triangle bounded by $x = 0$ , $x + y = 1$ , $y = 0$ .   |  |
| (h)        | C   | Find the flux of $F = vz \vec{i} + z^2 \vec{k}$ outward through the surface S cut from the  | 05   |
| (0)        | -   | cylinder $y^2 + z^2 = 1$ , $z \ge 0$ , by the planes $x = 0$ and $x = 1$  | 03   |
| (c)        |   | Cynnucl $y + z = 1, z \ge 0$ , by the planes $x = 0$ and $x = 1$ .<br>Use Stelse's theorem to explore $\int_{-\infty}^{\infty} E_{-x} dx$ if  |  |
|            |   | Use sloke sineorem to evaluate $\int_C r dr n$  | ••   |
|            |   | F = (x + y) i + (2x - z) i + (y + z) k and C is the boundary of the triangle  |  |
|            |   | (2,0,0), (0,3,0) and $(0,0,6)$ .  |  |
|            | <ul> <li>(b)</li> <li>(c)</li> <li>(a)</li> <li>(b)</li> <li>(c)</li> <li>(a)</li> <li>(b)</li> <li>(c)</li> <li>(c)</li> <li>(c)</li> <li>(c)</li> </ul> | <ul> <li>(b)</li> <li>(c)</li> <li>(a)</li> <li>(i)</li> <li>(i)</li></ul>  | (b) Find the equations for tangent plane and normal line at the point (1,1,1) on the surface $x^2 + y^2 + z^2 = 3$ .<br>(c) Find the area of the region that lies inside the cardioid $r = 1 + \cos \theta$ and outside the circle $r = 1$ .<br>OR<br>(a) Evaluate $\int_{0}^{1} \int_{0}^{\frac{1}{2}} \int_{0}^{2\pi} (r^2 \cos^2 \theta + z^2) r d\theta dr dz$ .<br>(b) (i) Integrate $f(x, y) = \frac{\ln(x^2 + y^2)}{\sqrt{x^2 + y^2}}$ over the region $1 \le x^2 + y^2 \le e$ by changing to polar coordinates<br>(ii) Find the derivative of $f(x, y, z) = x^3 - xy^2 - z$ at $P_0(1, 1, 0)$ in the direction of $\dot{v} = 2\dot{i} - 3\dot{j} + 6\dot{k}$ .<br>(c) Find the volume of the prism whose base is the triangle in $xy$ - plane bounded by the $x$ - axis and the line $y = x$ and $x = 1$ and whose top lies in the plane $z = f(x, y) = 3 - x - y$ .<br>(a) Integrate $f(x, y, z) = x + \sqrt{y} - z^2$ over the path $C_1 \cup C_2$ from $(0, 0, 0)$ to $(1, 1, 1)$ with $C_1 : r(t) = t\dot{i} + t\dot{j} + t\dot{k}$ , $0 \le t \le 1$<br>$C_2 : r(t) = \vec{i} + \vec{j} + t\dot{k}$ , $0 \le t \le 1$<br>(b) State Green's theorem and also evaluate the integral $\oint_C (y^2 dx + x^2 dy)$ where $f(x)$ the circle $(x - 2)^2 + (y - 3)^2 = 4$ .<br>(c) The triangle bounded by $x = 0$ , $x + y = 1$ , $y = 0$ .<br>(a) Use Green's theorem to evaluate the integral $\oint_C (y^2 dx + x^2 dy)$ where $f(x)$ the triangle bounded by $x = 0$ , $x + y = 1$ , $y = 0$ .<br>(b) Find the flux of $F = yz \vec{j} + z^2 \vec{k}$ outward through the surface S cut from the cylinder $y^2 + z^2 = 1$ , $z \ge 0$ , by the planes $x = 0$ and $x = 1$ .<br>(c) Use Stoke's theorem to evaluate $\int_C F \cdot dr$ if $F = (x + y) \vec{i} + (2x - z) \vec{j} + (y + z) \vec{k}$ and C is the boundary of the triangle $(2, 0, 0)$ (0, 3) and (0, 0, 6). |

\*\*\*\*\*

# Download all NOTES and PAPERS at StudentSuvidha.com