

B.E.(M.D.U.)  
First Semester Examination, 2009-10  
**Mathematics-1 (Math-1)**

**Note :** Attempt five questions in all, selecting two questions from each part.

**Part—(A)**

**Q. 1. (a) Test the convergence or divergence of the series :**

$$\sum \left[ \sqrt{n^4 + 1} - \sqrt{n^4 - 1} \right]$$

**Ans. Let**

$$\begin{aligned} u_n &= \sqrt{n^4 + 1} - \sqrt{n^4 - 1} \\ &= n^2 \sqrt{1 + \frac{1}{n^4}} - n^2 \sqrt{1 - \frac{1}{n^4}} \\ u_n &= n^2 \left[ \sqrt{1 + \frac{1}{n^4}} - \sqrt{1 - \frac{1}{n^4}} \right] \\ &= n^2 \left[ \left( 1 + \frac{1}{n^4} \right)^{1/2} - \left( 1 - \frac{1}{n^4} \right)^{1/2} \right] \\ &= n^2 \left\{ \left[ 1 + \frac{1}{2} \cdot \frac{1}{n^4} + \frac{\frac{1}{2} \left( \frac{1}{2} - 1 \right)}{2!} \frac{1}{n^8} + \frac{\frac{1}{2} \left( \frac{1}{2} - 1 \right) \left( \frac{1}{2} - 2 \right)}{3!} \frac{1}{n^{12}} + \dots \right] \right. \\ &\quad \left. - \left[ 1 - \frac{1}{2} \cdot \frac{1}{n^4} + \frac{\frac{1}{2} \left( \frac{1}{2} - 1 \right)}{2!} \frac{1}{n^8} - \dots \right] \right\} \\ &= n^2 \left[ 2 \left\{ \frac{1}{2n^4} + \frac{1}{16n^{12}} + \dots \right\} \right] \\ &= \frac{1}{n^2} + \frac{1}{8n^{10}} + \dots \end{aligned}$$

Let us take the auxiliary series  $\sum v_n = \sum \frac{1}{n^2}$

Now

$$\lim_{v_n} \frac{u_n}{v_n} = \lim \frac{\frac{1}{n^2} \left[ 1 + \frac{1}{8n^8} + \dots \right]}{\frac{1}{n^2}}$$

= 1 (which is finite and non-zero)

Since the auxiliary series  $\sum v_n = \sum \frac{1}{n^2}$  is convergent (as  $p = 2 > 1$ ). Hence by comparison test the given series  $\sum u_n$  is also convergent.

**Q. 1. (b) Discuss the convergence of the series**

$$x + \frac{2^2 x^2}{2!} + \frac{3^3 x^3}{3!} + \frac{4^4 x^4}{4!} + \dots \infty (x > 0)$$

**Ans. Let**

$$u_n = \frac{n^n x^n}{n!}$$

$$u_{n+1} = \frac{(n+1)^{n+1} x^{n+1}}{(n+1)!}$$

$$\frac{u_n}{u_{n+1}} = \frac{1}{x} \cdot \frac{n^n}{n!} \cdot \frac{(n+1)!}{(n+1)^{n+1}}$$

$$= \frac{1}{x} \cdot \frac{n^n (n+1)}{(n+1)^{n+1}}$$

$$= \frac{1}{x} \cdot \frac{n^n}{(n+1)^n}$$

$$= \frac{1}{x} \cdot \frac{1}{\left(1 + \frac{1}{n}\right)^n}$$

$$\lim_{n \rightarrow \infty} \frac{u_n}{u_{n+1}} = \frac{1}{x} \lim_{n \rightarrow \infty} \frac{1}{\left(1 + \frac{1}{n}\right)^n}$$

$$= \frac{1}{x} \cdot \frac{1}{e}$$

$$= \frac{1}{ex}$$

$\therefore$  By ratio test,  $\sum u_n$  is convergent if

$$\frac{1}{ex} > 1 \text{ i.e., } x < \frac{1}{e}$$

&  $\sum u_n$  is divergent if

$$\frac{1}{ex} < 1 \text{ i.e., } x > \frac{1}{e}$$

For  $\frac{1}{ex} = 1$ , i.e., for  $x = \frac{1}{e}$ , the ratio test fails.

Applying log test,

When

$$x = \frac{1}{e}$$

$$\frac{u_n}{u_{n+1}} = e \frac{1}{\left(1 + \frac{1}{n}\right)^n}$$

$$\begin{aligned} n \log \frac{u_n}{u_{n+1}} &= n \log \left[ e \frac{1}{\left(1 + \frac{1}{n}\right)^n} \right] \\ &= n \left[ \log e - \log \left(1 + \frac{1}{n}\right)^n \right] \\ &= n \left[ 1 - n \log \left(1 + \frac{1}{n}\right) \right] \\ &= n \left[ 1 - n \left( \frac{1}{n} - \frac{1}{2n^2} + \frac{1}{3n^3} - \dots \right) \right] \\ &= n \left( \frac{1}{2n} - \frac{1}{3n^2} + \dots \right) \\ &= \frac{1}{2} - \frac{1}{3n} + \dots \end{aligned}$$

$$\lim_{n \rightarrow \infty} n \log \left( \frac{u_n}{u_{n+1}} \right) = \frac{1}{2} < 1$$

By log test,  $\sum u_n$  is divergent for  $x = \frac{1}{e}$ . Hence the given series is convergent if  $x < \frac{1}{e}$  and is divergent if

$$x \geq \frac{1}{e}$$

**Q. 1. (c) State, with reasons, the values of  $x$  for which the series**

$$x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots \infty \text{ converges}$$

**Ans. Let**

$$\sum u_n = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

The series  $\sum u_n$  is absolutely convergent if the series  $\sum |u_n|$  is convergent. Applying ratio test,

$$\begin{aligned} \left| \frac{u_n}{u_{n+1}} \right| &= \left| \frac{x^n}{n} \cdot \frac{n+1}{x^{n+1}} \right| \\ &= \frac{n+1}{n} \cdot \frac{1}{|x|} \end{aligned}$$

$$= \left(1 + \frac{1}{n}\right) \frac{1}{|x|}$$

$$\lim \left| \frac{u_n}{u_{n+1}} \right| = \lim \left[ \left(1 + \frac{1}{n}\right) \frac{1}{|x|} \right] = \frac{1}{|x|}$$

So by ratio test, the series  $\sum |u_n|$  is convergent if  $\frac{1}{|x|} > 1$  i.e.,  $|x| < 1$  i.e.,  $-1 < x < 1$

$\therefore$  The given series is absolutely convergent and hence also convergent if  $-1 < x < 1$  if  $|x| < 1$

When  $x = 1$ , the given series is

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$$

Which converges by Leibnitz's test but converges conditionally.

**Q.2. (a) Compute to four decimal places, the value of  $\cos 32^\circ$ , by use of Taylor's series.**

**Ans.** To find  $\cos 32^\circ$        $\cos 32^\circ = \cos \frac{32\pi}{180}$

$$= \cos \frac{8\pi}{45}$$

Let us take  $\cos 32^\circ$  in the neighbours house of  $\cos 30^\circ$

$$\cos \left( \frac{8\pi}{45} + \frac{\pi}{6} - \frac{\pi}{6} \right) = \cos(x + h - h)$$

$$= f(x + h - h)$$

$$= f(a + h)$$

$$h = \left( \frac{8\pi}{45} - \frac{\pi}{6} \right), \quad a = \frac{\pi}{6}$$

$$h = \frac{48\pi - 45\pi}{270} = \frac{\pi}{90}$$

$$a = \frac{\pi}{6}$$

$$f(x) = \cos x \qquad f\left(\frac{\pi}{6}\right) = \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2}$$

$$f'(x) = -\sin x, \qquad f'\left(\frac{\pi}{6}\right) = -\sin \frac{\pi}{6} = -\frac{1}{2}$$

$$f''(x) = -\cos x, \qquad f''\left(\frac{\pi}{6}\right) = -\frac{\sqrt{3}}{2}$$

$$f'''(x) = \sin x, \qquad f'''\left(\frac{\pi}{6}\right) = \frac{1}{2}$$

$$f^{iv}(x) = \cos x, \quad f^{iv}\left(\frac{\pi}{6}\right) = \frac{\sqrt{3}}{2} \text{ and so on}$$

Now using Taylor's series  $f(x) = f(a) + hf'(a) + \frac{h^2}{2!} f''(a) + \dots$

$$\cos\left(\frac{8\pi}{45}\right) = \frac{\sqrt{3}}{2} + \frac{\pi}{90}\left(-\frac{1}{2}\right) + \frac{\pi^2}{90^2} \times \frac{1}{2!}\left(\frac{-\sqrt{3}}{2}\right) + \left(\frac{\pi}{90}\right)^3 \frac{1}{3!}\left(\frac{1}{2}\right) + \dots$$


$$\cos 32^\circ = \frac{\sqrt{3}}{2} - \frac{1}{2}(0.03492) - \frac{\sqrt{3}}{4}(0.00121945) + \frac{1}{12}(0.0000425832) + \dots$$

$$= 0.8660254 - 0.01746 - 0.00052802 + 0.0000035486$$

$$\cos 32^\circ = 0.84804 \text{ Ans.}$$

Q. 2. (b) If  $\rho$  be the radius of curvature at any point P on the parabola  $y^2 = 4ax$  and S be its focus, then show that  $\rho^2$  varies as  $(SP)^3$ .

Aus.



$$SP = \sqrt{(x-a)^2 + (y-0)^2}$$

$$(SP)^2 = (x-a)^2 + y^2$$

$$= x^2 + a^2 - 2ax + 4ax$$

$$[\because y^2 = 4ax]$$

$$(SP)^2 = (x+a)^2$$

$$SP = x+a$$

$$(SP)^3 = (x+a)^3$$

.... (i)

Now, given  $y^2 = 4ax$

Differentiating both sides w.r. to x,

$$2y \frac{dy}{dx} = 4a$$

$$\Rightarrow \frac{dy}{dx} = \frac{2a}{y} \quad \dots (ii)$$

$$1 + \left( \frac{dy}{dx} \right)^2 = 1 + \frac{4a^2}{y^2} = 1 + \frac{4a^2}{4ax}$$

$$1 + \left( \frac{dy}{dx} \right)^2 = \left( \frac{x+a}{x} \right)$$

Again differentiating equation (ii) w.r. to  $x$ ,

$$\frac{d^2 y}{dx^2} = -\frac{2a}{y^2} \frac{dy}{dx} = -\frac{2a}{4ax} \cdot \frac{2a}{y}$$

$$\frac{d^2 y}{dx^2} = -\frac{a}{xy}$$

Radius of curvature ( $\rho$ ) is given by

$$\rho = \frac{\left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^{3/2}}{\frac{d^2 y}{dx^2}}$$

$$\rho = \frac{\left( \frac{x+a}{x} \right)^{3/2}}{\left( -\frac{a}{xy} \right)}$$

$$\rho^2 = \left( \frac{x+a}{x} \right)^3 \times \frac{x^2 y^2}{a^2}$$

Or

$$\rho^2 = \left( \frac{x+a}{x} \right)^3 \times \frac{x^2 \cdot 4ax}{a^2}$$

$$\rho^2 = \frac{4}{a} (x+a)^3$$

$$\rho^2 = \frac{4}{a} (SP)^3 \quad [\text{from equation (i)}]$$

$$\rho^2 \propto (SP)^3 \quad \text{Hence proved}$$

Q. 2. (c) Show that the asymptotes of the curve  $x^2 y^2 = a^2 (x^2 + y^2)$  from a square of side  $2a$ .

Ans. Given curve is

$$x^2 y^2 = a^2 (x^2 + y^2)$$

For more study material Log on to <http://www.ululu.in/>

Since all powers of  $x$  and  $y$  are even asymptotes are parallel to  $x$  and  $y$ -axis.

**Asymptotes Parallel to  $x$ -axis :** Equating coefficient of highest power of  $x$  to zero i.e.,

$$y^2 - a^2 = 0$$


$y = \pm a$  are the asymptotes parallel to  $x$ -axis

**Asymptotes Parallel to  $y$ -axis :** By equating the coefficient of highest power of  $y$  to zero

$$x^2 - a^2 = 0$$

$x = \pm a$  are the asymptotes parallel to  $y$ -axis.

Since equation of the curve is of degree 4, it cannot have more than four asymptotes. Thus, four asymptotes are  $x = \pm a$ ,  $y = \pm a$ , which form a square



**Q. 3. (a) If  $u = x^2 \tan^{-1}(y/x) - y^2 \tan^{-1}(x/y)$ , evaluate**

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2}$$

**Ans. Given :**  $u = x^2 \tan^{-1}\left(\frac{y}{x}\right) - y^2 \tan^{-1}\left(\frac{x}{y}\right)$

Let  $u = v - w$

Where,  $v = x^2 \tan^{-1}\left(\frac{y}{x}\right)$ ,  $w = y^2 \tan^{-1}\left(\frac{x}{y}\right)$

$v$  is a homogeneous function of degree  $n = 2$  in  $x, y$ .

$$\begin{aligned} \text{Thus, } x^2 \frac{\partial^2 v}{\partial x^2} + 2xy \frac{\partial^2 v}{\partial x \partial y} + y^2 \frac{\partial^2 v}{\partial y^2} &= n(n-1)v \\ &= 2(2-1)v \\ &= 2v \end{aligned}$$

..... (i)

Also,  $w$  is also a homogeneous function of degree 2 in  $x, y$ .

$$x^2 \frac{\partial^2 w}{\partial x^2} + 2xy \frac{\partial^2 w}{\partial x \partial y} + y^2 \frac{\partial^2 w}{\partial y^2} = 2w$$

..... (ii)

Subtracting equation (ii) from equation (i)

$$\begin{aligned} x^2 \frac{\partial^2}{\partial x^2} (v-w) + 2xy \frac{\partial^2}{\partial x \partial y} (v-w) + y^2 \frac{\partial^2}{\partial y^2} (v-w) \\ = 2(v-w) \end{aligned}$$

$$\Rightarrow x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = 2u$$

Thus, we have

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = 2 \left\{ x^2 \tan^{-1} \left( \frac{y}{x} \right) - y^2 \tan^{-1} \left( \frac{x}{y} \right) \right\} \text{ Ans.}$$

**Q. 3. (b) If  $f(x, y) = \tan^{-1}(xy)$ , compute  $f(0.9, -1.2)$  approximately.**

**Ans.**  $f(x, y) = \tan^{-1}(xy)$

Let us expand  $f(x, y)$  near the point  $(1, -1)$

$$\begin{aligned} f(0.9, -1.2) &= f(1-0.1, -1-0.2) \\ &= f(1, -1) + \left[ (-0.1) \frac{\partial f}{\partial x} + (-0.2) \frac{\partial f}{\partial y} \right] \\ &\quad + \frac{1}{2!} \left[ (-0.1)^2 \frac{\partial^2 f}{\partial x^2} + 2(-0.1)(-0.2) \frac{\partial^2 f}{\partial x \partial y} + (-0.2)^2 \frac{\partial^2 f}{\partial y^2} \right] + \dots \dots \dots (i) \end{aligned}$$

Now,

$f(x, y) = \tan^{-1}(xy)$	$f(1, -1) = -\frac{\pi}{4}$
$\frac{\partial f}{\partial x} = \frac{y}{1+x^2y^2}$	$\frac{\partial f}{\partial x}(1, -1) = -\frac{1}{2}$
$\frac{\partial f}{\partial y} = \frac{x}{1+x^2y^2}$	$\frac{\partial f}{\partial y}(1, -1) = \frac{1}{2}$
$\frac{\partial^2 f}{\partial x^2} = \frac{-2xy}{(1+x^2y^2)^2}$	$\frac{\partial^2 f}{\partial x^2}(1, -1) = \frac{1}{2}$
$\frac{\partial^2 f}{\partial x \partial y} = \frac{1+x^2y^2-2x^2y^2}{(1+x^2y^2)^2}$	$\frac{\partial^2 f}{\partial x \partial y}(1, -1) = 0$
$\frac{\partial^2 f}{\partial y^2} = \frac{-x(2x^2y)}{(1+x^2y^2)^2}$	$\frac{\partial^2 f}{\partial y^2}(1, -1) = \frac{1}{2}$

Putting all these values in equation (i),

$$\begin{aligned} f(0.9, -1.2) &= -\frac{\pi}{4} + (-0.1) \left( -\frac{1}{2} \right) + (-0.2) \left( \frac{1}{2} \right) \\ &\quad + \frac{1}{2} \left[ (-0.1)^2 \left( \frac{1}{2} \right) + 2(-0.1)(-0.2)(0) + (-0.2)^2 \left( \frac{1}{2} \right) \right] + \dots \dots \dots \\ &= -\frac{\pi}{4} + 0.05 - 0.1 + \frac{1}{2} (0.005 + 0.02) \\ &= -\frac{\pi}{4} + 0.05 - 0.1 + 0.0125 \\ f(0.9, -1.2) &= -0.823 \text{ Ans.} \end{aligned}$$



**Q. 4. (a) Find the maximum and minimum distances from the origin to the curve**  
 $5x^2 + 6xy + 5y^2 - 8 = 0$

**Ans.** Let  $p(x, y)$  be any point on the curve. Distance of the point  $A(0, 0)$  from  $P(x, y)$  is

$$\sqrt{(x-0)^2 + (y-0)^2}$$

If the distance is maximum or minimum, so will be the square of the distance.

Let  $f(x, y) = x^2 + y^2$  ..... (i)

Subject to the condition  $\phi(x, y) = 5x^2 + 6xy + 5y^2 - 8 = 0$  ..... (ii)

Consider Lagrange's function

$$F(x, y) = f(x, y) + \lambda \phi(x, y)$$

$$F(x, y) = x^2 + y^2 + \lambda(5x^2 + 6xy + 5y^2 - 8)$$

For stationary values  $dF = 0$

$$\{2x + \lambda(10x + 6y)\}dx + \{2y + \lambda(6x + 10y)\}dy = 0$$

$$2x + \lambda(10x + 6y) = 0$$

$$2y + \lambda(6x + 10y) = 0$$

Multiplying equation (iii) by  $x$  and equation (iv) by  $y$  and on adding

$$2(x^2 + y^2) + \lambda(10x^2 + 6xy + 6xy + 10y^2) = 0$$

$$2(x^2 + y^2) + 2\lambda(5x^2 + 6xy + 5y^2) = 0$$

$$\Rightarrow f(x, y) + \lambda(8) = 0$$

$$\lambda = -\frac{f}{8} \quad (\text{using equations (i) and (ii)})$$

From equations (iii) and (iv)

$$2x - \frac{f}{8}(10x + 6y) = 0$$

$$2y - \frac{f}{8}(6x + 10y) = 0$$

$$\Rightarrow 4x - f(5x + 3y) = 0$$

$$4y - f(3x + 5y) = 0$$

Or  $(4 - 5f)x - 3fy = 0$  .... (v)

$$-3fx + (4 - 5f)y = 0$$
 ..... (vi)

Solving equations (v) and (vi)

$$(3f)^2 = (4 - 5f)^2$$

$$9f^2 = 16 + 25f^2 - 40f$$

$$16f^2 - 40f + 16 = 0$$

$$2f^2 - 5f + 2 = 0$$

$$2f^2 - 4f - f + 2 = 0$$

$$2f(f-2) - 1(f-2) = 0$$

$$(2f-1)(f-2) = 0$$

$$f = \frac{1}{2}, 2$$

Thus, the maximum distance  $= \sqrt{2}$   
 $= 1.414$

& minimum distance  $= \sqrt{\frac{1}{2}}$   
 $= 0.7072$  Ans.

Q. 4. (b) Evaluate

$$\int_0^{\alpha} \frac{\log(1+\alpha x)}{1+x^2} dx$$

Ans.

$$\int_0^{\alpha} \frac{\log(1+\alpha x)}{1+x^2} dx$$

Let us take

$$\alpha = 1$$

$$I = \int_0^1 \frac{\log(1+x)}{1+x^2} dx$$

Now putting

$$x = \tan \theta$$

$$dx = \sec^2 \theta d\theta$$

When  $x = 0$ ,  $\theta = 0$

When

$$x = 1, \theta = \frac{\pi}{4}$$

$$I = \int_0^{\pi/4} \frac{\log(1+\tan \theta)}{(1+\tan^2 \theta)} \sec^2 \theta d\theta$$

$$I = \int_0^{\pi/4} \log(1+\tan \theta) d\theta$$

Applying property equation (iv) of definite integral

$$I = \int_0^{\pi/4} \log \left\{ 1 + \tan \left( \frac{\pi}{4} - \theta \right) \right\} d\theta$$

$$\begin{aligned}
&= \int_0^{\pi/4} \log \left\{ 1 + \left( \frac{\tan \frac{\pi}{4} - \tan \theta}{1 + \tan \frac{\pi}{4} \tan \theta} \right) \right\} d\theta \\
&= \int_0^{\pi/4} \log \left( 1 + \frac{1 - \tan \theta}{1 + \tan \theta} \right) d\theta \\
&= \int_0^{\pi/4} \log \left( \frac{2}{1 + \tan \theta} \right) d\theta \\
I &= \int_0^{\pi/4} \log 2 \, d\theta - \int_0^{\pi/4} \log(1 + \tan \theta) d\theta \\
I &= \int_0^{\pi/4} \log 2 \, d\theta - I \\
2I &= \int_0^{\pi/4} \log 2 \, d\theta \\
&= \log 2 \int_0^{\pi/4} 1 \, d\theta \\
2I &= \log 2 [\theta]_0^{\pi/4} \\
2I &= \frac{\pi}{4} \log 2 \\
I &= \frac{\pi}{8} \log 2 \quad \text{Ans.}
\end{aligned}$$

#### Part—(B)

**Q. 5. (a)** Find the volume of solid formed by revolving a loop of the lemniscate  $r^2 = a^2 \cos 2\theta$  about the initial line.

**Ans.** For the upper half of the loop  $\theta$  varies from 0 to  $\pi/4$ . The curve is revolving about the initial line i.e., x-axis

$$\begin{aligned}
\text{Required volume} &= \frac{2}{3} \pi \int_0^{\pi/4} r^3 \sin \theta \, d\theta \\
&= \frac{2}{3} \pi \int_0^{\pi/4} \{a\sqrt{\cos 2\theta}\}^3 \sin \theta \, d\theta && [\because r^2 = a^2 \cos 2\theta] \\
&= \frac{2\pi a^3}{3} \int_0^{\pi/4} (2\cos^2 \theta - 1)^{3/2} \sin \theta \, d\theta
\end{aligned}$$

Put

$$\sqrt{2} \cos \theta = \sec \phi$$

$$-\sqrt{2} \sin \theta d\theta = \sec \phi \tan \phi d\phi$$

& when  $\theta = 0$ ,  $\phi = \pi/4$  and when  $\theta = \frac{\pi}{4}$ ,  $\phi = 0$

$$= \frac{2\pi a^3}{3} \int_{\pi/4}^0 (\sec^2 \theta - 1)^{3/2} \frac{(-\sec \phi \tan \phi)}{\sqrt{2}} d\phi$$

$$= \frac{\sqrt{2}\pi a^3}{3} \int_0^{\pi/4} \tan^4 \phi \sec \phi d\phi$$

$$= \frac{\sqrt{2}\pi a^3}{3} \int_0^{\pi/4} (\sec^2 \phi - 1)^2 \sec \phi d\phi$$

$$= \frac{\sqrt{2}\pi a^3}{3} \int_0^{\pi/4} (\sec^5 \phi - 2\sec^3 \phi + \sec \phi) d\phi \quad \dots (i)$$

Now using the reduction formula

$$\int \sec^n \phi d\phi = \frac{\sec^{n-2} \phi \tan \phi}{n-1} + \frac{(n-2)}{(n-1)} \int \sec^{n-2} \phi d\phi$$

Thus,

$$\int_0^{\pi/4} \sec^5 \phi d\phi = \left[ \frac{\sec^3 \phi \tan \phi}{4} \right]_0^{\pi/4} + \frac{3}{4} \int_0^{\pi/4} \sec^3 \phi d\phi$$

$$= \frac{\sqrt{2}}{2} + \frac{3}{4} \left[ \left\{ \frac{\sec \phi \tan \phi}{2} \right\}_0^{\pi/4} + \frac{1}{2} \int_0^{\pi/4} \sec \phi d\phi \right]$$

$$= \frac{\sqrt{2}}{2} + \frac{3}{4} \left[ \frac{\sqrt{2}}{2} + \frac{1}{2} \{ \log(\sec \phi + \tan \phi) \}_0^{\pi/4} \right]$$

$$= \frac{\sqrt{2}}{2} + \frac{3\sqrt{2}}{8} + \frac{3}{8} \log(\sqrt{2} + 1) = \frac{7\sqrt{2}}{8} + \frac{3}{8} \log(\sqrt{2} + 1)$$

&

$$\int_0^{\pi/4} \sec^3 \phi d\phi = \left[ \frac{\sec \phi \tan \phi}{2} \right]_0^{\pi/4} + \frac{1}{2} \int_0^{\pi/4} \sec \phi d\phi$$

$$= \frac{\sqrt{2}}{2} + \frac{1}{2} \log(\sqrt{2} + 1)$$

&

$$\int_0^{\pi/4} \sec \phi d\phi = \log(\sqrt{2} + 1)$$


From equation (i), Required Volume

$$\begin{aligned}
 &= \frac{\sqrt{2}\pi a^3}{3} \left[ \frac{7\sqrt{2}}{8} + \frac{3}{8} \log(\sqrt{2}+1) - 2 \left\{ \frac{\sqrt{2}}{2} + \frac{1}{2} \log(\sqrt{2}+1) \right\} + \log(\sqrt{2}+1) \right] \\
 &= \frac{\sqrt{2}\pi a^3}{3} \left[ \frac{3}{8} \log(\sqrt{2}+1) - \frac{\sqrt{2}}{8} \right] \\
 V &= \frac{\pi a^3 \sqrt{2}}{24} [3 \log(\sqrt{2}+1) - \sqrt{2}] \quad \text{Ans.}
 \end{aligned}$$

Q. 5. (b) Evaluate  $\int_0^{4a} \int_{x^2/4a}^{2\sqrt{ax}} dy dx$  by changing the order of integration.

Ans. Let

$$I = \int_0^{4a} \int_{x^2/4a}^{2\sqrt{ax}} dy dx$$



By changing the order of integration

$$\begin{aligned}
 I &= \int_0^{4a} \int_{y^2/4a}^{2\sqrt{ay}} dx dy \\
 I &= \int_0^{4a} [x]_{y^2/4a}^{2\sqrt{ay}} dy \\
 &= \int_0^{4a} \left[ 2\sqrt{ay} - \frac{y^2}{4a} \right] dy \\
 &= 2\sqrt{a} \left[ \frac{2}{3} y^{3/2} \right]_0^{4a} - \frac{1}{4a} \left[ \frac{y^3}{3} \right]_0^{4a}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{4\sqrt{a}}{3}(4a)^{3/2} - \frac{1}{12a}(4a)^3 \\
&= \frac{32a^2}{3} - \frac{16a^2}{3} \\
&= \frac{16a^2}{3} \quad \text{Ans.}
\end{aligned}$$

Q. 6. (a) Evaluate  $\iiint (x+y+z) dx dy dz$  over the tetrahedron bounded by the planes  $x=0$ ,  $y=0$ ,  $z=0$  and  $x+y+z=1$ .

Ans. To evaluate  $\iiint (x+y+z) dx dy dz$  over the tetrahedron bounded by the planes  $x=0$ ,  $y=0$ ,  $z=0$  and  $x+y+z=1$

$$\begin{aligned}
&\iiint (x+y+z) dx dy dz \\
&= \iiint x^{1-1} y^{1-1} z^{1-1} (x+y+z) dx dy dz
\end{aligned}$$

Where  $0 \leq x+y+z \leq 1$

$$\begin{aligned}
\text{Using Liouville's extension} \quad &= \frac{1! 1! 1!}{1+1+1!} \int_0^1 u \cdot u^{1+1+1-1} \cdot du \\
&= \frac{1}{3!} \int_0^1 u \cdot u^2 du \\
&= \frac{1}{2} \int_0^1 u^3 du \\
&= \frac{1}{2} \left[ \frac{u^4}{4} \right]_0^1 \\
&= \frac{1}{8} [u^4]_0^1 \\
&= \frac{1}{8} \quad \text{Ans.}
\end{aligned}$$

Q. 6. (b) Show that

$$\int_0^{\pi/2} \sqrt{\sin \theta} d\theta \times \int_0^{\pi/2} \frac{d\theta}{\sqrt{\sin \theta}} = \pi$$

Ans. Taking L.H.S.

$$\int_0^{\pi/2} \sqrt{\sin \theta} d\theta \times \int_0^{\pi/2} \frac{d\theta}{\sqrt{\sin \theta}}$$

⇒

$$\int_0^{\pi/2} \sin^{1/2} \theta \cos^0 \theta d\theta \times \int_0^{\pi/2} \sin^{-1/2} \theta \cos^0 \theta d\theta$$

$$\left[ \because \int_0^{\pi/2} \sin^p \theta \cos^q \theta d\theta = \frac{\frac{p+1}{2} \frac{q+1}{2}}{2 \frac{p+q+2}{2}} \right]$$

⇒

$$\frac{\frac{\frac{1}{2}+1}{2} \frac{0+1}{2}}{2 \frac{\frac{1}{2}+0+2}{2}} \times \frac{\frac{\frac{-1}{2}+1}{2} \frac{0+1}{2}}{2 \frac{\frac{-1}{2}+0+2}{2}}$$

$$= \frac{\frac{\frac{3}{4}}{2} \frac{\frac{1}{2}}{2}}{2 \frac{\frac{5}{4}}{2}} \times \frac{\frac{\frac{1}{4}}{2} \frac{\frac{1}{2}}{2}}{2 \frac{\frac{3}{4}}{2}}$$

$$= \frac{\left(\frac{1}{2}\right)^2 \frac{1}{4}}{4 \frac{5}{4}}$$

$$= \frac{(\sqrt{\pi})^2 \frac{1}{4}}{4 \times \frac{1}{4} \frac{1}{4}}$$

$$= (\sqrt{\pi})^2$$

$$= \pi = \text{R.H.S.}$$

$$\left[ \because \frac{1}{2} = \sqrt{\pi} \quad \& \quad n! = (n-1)n-1! \right]$$

**Q. 7. (a) Find the constants  $a$  and  $b$  so that the surface  $ax^2 - byz = (a+2)x$  is orthogonal to the surface  $4x^2y + z^3 = 4$  at the point  $(1, -1, 2)$ .**

Ans. Given the two surfaces are

$$ax^2 - byz = (a+2)x$$

&

$$4x^2y + z^3 = 4$$

Let

$$\phi_1 \equiv ax^2 - byz - (a+2)x \quad \dots (i)$$

$$\phi_2 \equiv 4x^2y + z^3 - 4 \quad \dots (ii)$$

$$\nabla \phi_1 \equiv \left( \hat{i} \frac{\partial}{\partial x} + \hat{j} \frac{\partial}{\partial y} + \hat{k} \frac{\partial}{\partial z} \right) (ax^2 - byz - ax - 2x)$$

$$= (2ax - (a+2))\hat{i} + (-bz)\hat{j} + (-by)\hat{k}$$

At (1, -1, 2)

$$\nabla\phi_1 = (a-2)\hat{i} - 2b\hat{j} + b\hat{k} \quad \dots (iii)$$

Also

$$\nabla\phi_2 = \left( \hat{i} \frac{\partial}{\partial x} + \hat{j} \frac{\partial}{\partial y} + \hat{k} \frac{\partial}{\partial z} \right) (4x^2y + z^3 - 4)$$

$$\nabla\phi_2 = 8xy\hat{i} + 4x^2\hat{j} + 3z^2\hat{k}$$

At (1, -1, 2)

$$\nabla\phi_2 = -8\hat{i} + 4\hat{j} + 12\hat{k} \quad \dots (iv)$$

Since both surfaces are orthogonal,

$$\nabla\phi_1 \cdot \nabla\phi_2 = 0$$

$$\{(a-2)\hat{i} - 2b\hat{j} + b\hat{k}\} \cdot \{-8\hat{i} + 4\hat{j} + 12\hat{k}\} = 0$$

$$\Rightarrow -8(a-2) - 8b + 12b = 0$$

$$-8a + 4b + 16 = 0 \quad \dots (v)$$

Also since both surfaces are orthogonal at (1, -1, 2) so this point will satisfy the surfaces. Thus, from first surface

$$a(1)^2 - b(-1)(2) = (a+2)(1)$$

$$a + 2b - a - 2 = 0$$

$$2b = 2$$

$$b = 1$$

Putting  $b = 1$  in equation (v)

$$-8a + 4(1) + 16 = 0$$

$$-8a = -20$$

$$a = \frac{20}{8}$$

$$a = \frac{5}{2}$$

The values of  $a$  and  $b$  are  $\frac{5}{2}$  and 1.

**Q. 7. (b)** If  $v_1$  and  $v_2$  be the vectors joining the fixed points  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  respectively to a variable point  $(x, y, z)$ , prove that

$$\text{curl}(v_1 \times v_2) = 2(v_1 - v_2)$$

**Ans.**

$$\vec{V}_1 = x_1\hat{i} + y_1\hat{j} + z_1\hat{k}$$

$$\vec{V}_2 = x_2\hat{i} + y_2\hat{j} + z_2\hat{k}$$




$$\begin{aligned}\vec{V}_1 \times \vec{V}_2 &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \end{vmatrix} \\ &= \hat{i}(y_1 z_2 - y_2 z_1) - \hat{j}(x_1 z_2 - x_2 z_1) + \hat{k}(x_1 y_2 - x_2 y_1) \\ \vec{V}_1 \times \vec{V}_2 &= \hat{i}(y_1 z_2 - y_2 z_1) + \hat{j}(x_2 z_1 - x_1 z_2) + \hat{k}(x_1 y_2 - x_2 y_1)\end{aligned}$$

Now curl  $(\vec{V}_1 \times \vec{V}_2) = \nabla \times (\vec{V}_1 \times \vec{V}_2)$

$$\begin{aligned}&= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ (y_1 z_2 - y_2 z_1) & (x_2 z_1 - x_1 z_2) & (x_1 y_2 - x_2 y_1) \end{vmatrix} \\ &= \left[ \frac{\partial}{\partial y} (x_1 y_2 - x_2 y_1) - \frac{\partial}{\partial z} (x_2 z_1 - x_1 z_2) \right] \hat{i} - \left[ \frac{\partial}{\partial x} (x_1 y_2 - x_2 y_1) - \frac{\partial}{\partial z} (y_1 z_2 - y_2 z_1) \right] \hat{j} \\ &\quad + \hat{k} \left[ \frac{\partial}{\partial x} (x_2 z_1 - x_1 z_2) - \frac{\partial}{\partial y} (y_1 z_2 - y_2 z_1) \right] \\ &= \hat{i}[(x_1 - x_2) - (x_2 - x_1)] - \hat{j}[(y_2 - y_1) - (y_1 - y_2)] + \hat{k}[(z_1 - z_2) - (z_2 - z_1)] \\ &= \hat{i}(2x_1 - 2x_2) - \hat{j}(2y_2 - 2y_1) + \hat{k}(2z_1 - 2z_2) \\ &= 2[\hat{i}(x_1 - x_2) + \hat{j}(y_1 - y_2) + \hat{k}(z_1 - z_2)] \\ &= 2[(x_1 \hat{i} + y_1 \hat{j} + z_1 \hat{k}) - (x_2 \hat{i} + y_2 \hat{j} + z_2 \hat{k})] \\ &= 2(\vec{V}_1 - \vec{V}_2) \\ &= \text{R.H.S.}\end{aligned}$$

**Q. 8. (a) Verify Stoke's theorem for  $F = (x^2 + y^2)\hat{i} - 2xy\hat{j}$  taken around the rectangle bounded by the lines  $x = \pm a$ ,  $y = 0$ ,  $y = b$ .**

**Ans.** Let C denote the boundary of the rectangle ABED, then



$$\oint_C \vec{F} \cdot d\vec{r} = \oint_C [(x^2 + y^2)\hat{i} - 2xy\hat{j}] \cdot (\hat{i}dx + \hat{j}dy)$$

$$= \oint_C (x^2 + y^2)dx - 2xydy$$

The curve  $C$  consists of four lines  $AB$ ,  $BE$ ,  $ED$  and  $DA$ .

Along  $AB$ ,  $x = a$ ,  $dx = 0$ ,  $y$  varies from  $0$  to  $b$

$$\int_{AB} (x^2 + y^2)dx - 2xydy$$

$$= \int_0^b -2aydy = -2a \left[ \frac{y^2}{2} \right]_0^b = -ab^2$$

.... (i)

Along  $BE$ ,  $y = b$ ,  $dy = 0$ ,  $x$  varies from  $a$  to  $-a$

$$\int_{BE} (x^2 + y^2)dx - 2xydy = \int_a^{-a} (x^2 + b^2)dx$$

$$= \left[ \frac{x^3}{3} + b^2x \right]_a^{-a}$$

$$= \frac{-a^3}{3} - ab^2 - \frac{a^3}{3} - ab^2$$

$$= \frac{-2a^3}{3} - 2ab^2$$

.... (ii)

Along  $ED$ ,  $x = -a$ ,  $dx = 0$ ,  $y$  varies from  $b$  to  $0$

$$\int_{ED} (x^2 + y^2)dx - 2xydy = \int_b^0 2aydy = -ab^2$$

.... (iii)

Along  $DA$ ,  $y = 0$ ,  $dy = 0$ ,  $x$  varies from  $-a$  to  $a$

$$\int_{DA} (x^2 + y^2)dx - 2xydy = \int_{-a}^a x^2dx = \frac{2a^3}{3}$$

..... (iv)

On adding equations (i), (ii), (iii) and (iv)

$$\oint_C \vec{F} \cdot d\vec{r} = -ab^2 - \frac{2a^3}{3} - 2ab^2 - ab^2 + \frac{2a^3}{3}$$

$$= -4ab^2$$

.... (v)

Now  $\text{curl } \vec{F} = \nabla \times \vec{F}$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ (x^2 + y^2) & -2xy & 0 \end{vmatrix}$$

$$= (-2y - 2y)\hat{k} = -4y\hat{k}$$

For the surface  $S$ ,  $\hat{n} = \hat{k}$

$$\Rightarrow \text{curl } \vec{F} \cdot \hat{n} = -4y\hat{k} \cdot \hat{k} = -4y$$

Now 
$$\iint_S \text{curl } \vec{F} \cdot \hat{n} dS = \int_0^b \int_a^a -4y dx dy$$

$$= \int_0^b -4y [x]_{-a}^a dy$$

$$= -8a \int_0^b y dy$$

$$= -4a [y^2]_0^b$$

$$= -4ab^2$$

.... (vi)

From equations (v) and (vi)

$$\oint_C \vec{F} \cdot d\vec{r} = \iint_S \text{curl } \vec{F} \cdot \hat{n} dS$$

Hence verifies Stoke's theorem.

**Q. 8. (b) Using divergence theorem, evaluate  $\int_S \vec{R} \cdot \hat{n} ds$  where  $S$  is the surface of the sphere**

$$x^2 + y^2 + z^2 = 9.$$

Ans. To evaluate 
$$\iint_S \vec{r} \cdot \hat{n} dS$$

Using divergence theorem,

$$\iint_S \vec{r} \cdot \hat{n} dS = \iiint_V \text{div } \vec{r} dV$$

... (i)

We know that 
$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\text{div } \vec{r} = \nabla \cdot \vec{r}$$

$$= \left( \hat{i} \frac{\partial}{\partial x} + \hat{j} \frac{\partial}{\partial y} + \hat{k} \frac{\partial}{\partial z} \right) (x\hat{i} + y\hat{j} + z\hat{k})$$

$$= \frac{\partial}{\partial x} (x) + \frac{\partial}{\partial y} (y) + \frac{\partial}{\partial z} (z)$$

$$= 1 + 1 + 1$$

$$= 3$$

From equation (i)  $\iint_S \hat{n} dS = \iiint_V 3 dV$

$$= 3 \iiint_V dV$$

$$= 3 \text{ Volume of the given sphere with radius 3}$$

$$= 3 \cdot \frac{4}{3} \pi r^3$$

$$= 4\pi r^3$$

$$= 4\pi(3)^3$$

$$\iint_S \hat{n} dS = 108\pi \quad \text{Ans.}$$