

Name of the Course: **LOCF B.Sc. (H) Mathematics**
 Unique Paper Code: **32357501**
 Name of the Paper: **DSE-1 (i) Numerical Analysis**
 Semester: **V**
 Duration: **3 hours**
 Maximum Marks: **75 Marks**

Attempt any four questions. All questions carry equal marks.

1. (a) Find the smallest positive root of the given equation by performing three iterations of the Bisection method

$$f(x) = e^x - 3x = 0.$$

- (b) Apply four iterations of the Fixed Point Iteration Method to find an approximate root of the following equation by taking the initial approximation as $p_0=0$

$$f(x) = 3x - (1 + \cos x) = 0.$$

Also represent the root graphically.

- (c) Find LU decomposition for the matrix

$$A = \begin{pmatrix} 3 & 1 & 2 \\ 2 & -3 & -1 \\ 1 & -2 & -1 \end{pmatrix}$$

and then solve the system

$$3x + y + 2z = 4$$

$$2x - 3y - z = -6$$

$$x - 2y - z = -4$$

2. (a) Set up the Gauss-Seidel iteration scheme for the following system of equations

$$2x - y + 2z = 3$$

$$x + 3y + 3z = -1$$

$$x + 2y + 5z = 1$$

and iterate three times starting with the initial vector $X^{(0)} = (0,0,0.5)^T$.

- (b) Apply Secant method to find a root of the equation

$$\ln(1+x) - \frac{1}{2} \cos x = 0$$

(0,1). Perform three iterations. What order of convergence do you expect?

- (c) Solve the following system of equations using SOR iteration method

$$5x_1 + x_2 - 2x_3 = 2$$

$$3x_1 + 4x_2 - x_3 = -2$$

$$2x_1 - 3x_2 + 5x_3 = 10$$

Take $w = 0.9$ with $X^{(0)} = (0,0,0)^T$ and iterate three times.

3. (a) Find a polynomial of degree 3 or less passing through the points $(-1,9)$, $(0,5)$, $(1,3)$ and $(2,1)$ using Lagrange interpolation. Use this polynomial to estimate the ordinate for $x = 1.5$.
- (b) Use central difference formula to approximate $f'(0.5)$ for the function $f(x) = 4e^{-2x}$ by taking two step sizes $h = 0.1$ and 0.05 . What is the order of approximation?

4. (a) Derive the following approximation of function $f'(x_0)$ for an arbitrary function $f(x)$:

$$f'(x_0) = \frac{-3f_0 + 4f_1 - f_2}{2h}$$

Hence show that the above approximation provides exact value of the first order derivatives for $f(x) = 1, x$ and x^2 but not for $f(x) = x^3$.

(b) Determine the step size h in an equidistant table for $f(x) = \sin x$ in $[0, \pi/4]$, if the error in magnitude in quadratic interpolation is less than or equal to 5×10^{-8} in magnitude.

(c) Obtain the piecewise linear interpolating polynomials for the function $f(x)$ defined by the given data .

X	0.5	1.5	2.5
f(x)	0.125	3.375	15.625

Interpolate at $x = 1.0$ and 2.0 .

5. (a) Approximate the value of the given integral by Simpson's 1/3 rule and the Trapezoidal rule

$$\int_0^3 \frac{1}{4+x^2} dx.$$

Calculate the difference between the actual value and the approximate value. Also find the error term for both.

(b) Apply the Newton Raphson Method to find a root of the equation

$$f(x) = x^3 - x^2 - 10x + 6 = 0. \text{ Perform three iterations with } x_0 = 1.$$

- 6 (a) Construct the Richardson extrapolation table to find the derivative of the function $f(x) = \tan^{-1} x$ at $x_0 = 3$ using the first-order forward difference approximation

$$D_h^{(1)} = \frac{f(x_0 + h) - f(x_0)}{h}$$

taking $h=1,0.5,0.25,0.125$.

(b) Apply the Modified Euler method to find approximate solution of the following initial value problem with four numbers of steps

$$\frac{dy}{dx} = xy, \quad (1 \leq x \leq 2), \quad y(1) = 1.$$

Also compare with the exact solution.

downloaded from
StudentSuvidha.com