

Name of Course	: <b>CBCS B.Sc. (H) Mathematics</b>
Unique Paper Code	: <b>32351601</b>
Name of Paper	: <b>C 13- Complex Analysis</b>
Semester	: <b>VI</b>
Duration	: <b>3 hours</b>
Maximum Marks	: <b>75 Marks</b>

*Attempt any four questions. All questions carry equal marks.*

1. Determine whether  $S = \{z \in \mathbb{C} : |z|^2 > z + \bar{z}\}$  is a domain or not? Justify your answer.

Find the image of line segment joining  $z_1 = -i$  to  $z_2 = -1$  under the map  $f(z) = \bar{iz}$ .

Check whether Cauchy-Riemann equations for  $f(z) = \sqrt{|z^2 - \bar{z}^2|}$  are satisfied at the origin? Is  $f$  analytic at the origin? Justify your answer.

Suppose  $f(z) = \cosh(2x) \cos(2y) + i v(x, y)$  is analytic everywhere such that  $v(0,0) = 0$ . Find  $f(z)$ . Hence find zeros of  $f$ .

Solve the equation  $e^{z-1} + ie^3 = 0$ .

2. Let  $S = \{z \in \mathbb{C} : \text{Im } z = 1 \text{ and } \text{Re } z \neq 4\}$ . Is  $S$  open? Is  $S$  closed? Justify your answer.

Assume that  $g$  is analytic in a region and that at every point either  $g = 0$  or  $g' = 0$ . Show that  $g$  is constant.

Suppose  $f(z) = \begin{cases} \bar{z}^3/z^2 & \text{if } z \neq 0 \\ 0 & \text{if } z = 0 \end{cases}$ . Show that  $f$  is continuous everywhere on  $\mathbb{C}$ . Is  $f$  analytic at  $z = 0$ ? Justify your answer.

Does there exist an analytic function  $f(z) = u(x, y) + i v(x, y)$  for which  $u(x, y) = y^3 + 5x$ ? Solve the equation  $\text{Log}(z) + \text{Log}(2z) = 3\pi/2$ .

3. Determine whether the following curves are simple, closed, smooth or contour

$$C_1: z(t) = |t| + it, \quad t \in [-1, 1]$$

$$C_2: z(t) = e^{2it}, \quad t \in [0, 2\pi],$$

$C_3: z(t)$  is the positively oriented boundary of the rectangle whose sides lie along  $x = \pm 1, y = 0, y = 1$ .

Evaluate  $\int_{C_3} |z| dz$ . Explain why Cauchy Goursat theorem is not applicable in this case?

Use ML-Inequality to show that

$$\left| \int_C \frac{e^z}{(z+1)} dz \right| \leq 4\pi e^2$$

where  $C : z(t) = e^{2it}, t \in [-\pi, \pi]$ .

4. Evaluate  $\int_C ze^{3z} dz$  where  $C$  is the parabola  $x^2 = y$  from  $(0,0)$  to  $(1,1)$ .

Using Cauchy Integral formula, determine the integral  $\int_C \frac{e^z}{z^2(z^2-9)} dz$  where  $C$  is positively oriented circle (i)  $C : |z| = 1$ . (ii)  $C : |z - 3| = 1$ .

Use Liouville's theorem to establish that  $\cos z$  is not bounded in the complex plane.

Let  $g$  be an entire function and suppose that  $|g(z)| < 10$  for all values of  $z$  on the circle  $|z - 2| = 3$ . Find a bound for  $|g'''(2)|$ .

5. Determine the radius of convergence of the series  $\sum_{k=0}^{\infty} \frac{z^k}{k!}$  and  $\sum_{k=0}^{\infty} k^k z^k$ . Also discuss the convergence of the series.

Obtain the Maclaurin series of the function  $f(z) = \frac{1}{z^2} \sinh\left(\frac{1}{z}\right)$ . Specify the region in which the series is valid.

Find the Laurent series of the function  $f(z) = \frac{1}{(z+1)(z+3)}$  valid for  $0 < |z + 1| < 2$ .

6. Determine the residue and singularities of the function  $g(z) = \frac{z+1}{z^2+4}$ . Also evaluate  $\int_C g(z) dz$  where  $C$  is the positively oriented circle  $|z - i| = 2$ .

Using a single residue, evaluate the integral  $\int_{C'} \frac{3z-1}{z(z+1)} dz$  where  $C'$  is the positively oriented circle  $|z - 1| = 4$ .

Use residue to evaluate the integral  $\int_0^{2\pi} \frac{dt}{3+\cos t}$

downloaded from  
StudentSuvidha.com