Roll	No.	******************************
		*********************

## 21255

# B. Sc. (Hons.) Mathematics 2nd Semester Examination – May, 2019

## REGRESSION ANALYSIS AND PROBABILITY

Paper: BHM-125 Opt-i

Time: Three hours ]

[ Maximum Marks : 60

Before answering the questions candidates should ensure that they have been supplied the correct and complete question paper. No complaint whis regard will be entertained after examination.

Note: Attempt five questions in all, selecting one question from each Unit. Question No. 9 is compulsory.

### UNIT -

(a) A student while calculating the coefficient of correlation between two variates X and Y from 25 pairs of observations obtained the following values N = 25, ΣX = 125, ΣY = 100, ΣX<sup>2</sup> = 650 ΣY<sup>2</sup> = 460, ΣXY = 508. It was, detected later on the time of checking that he had copied down two

pairs of observations are  $\begin{bmatrix} X & Y \\ 6 & 14 \end{bmatrix}$  while the correct values are  $\begin{bmatrix} X & Y \\ 8 & 6 \end{bmatrix}$  values are  $\begin{bmatrix} X & Y \\ 8 & 12 \end{bmatrix}$  btain the correct value of the coefficient of correlation.

(b) Obtain the two lines of regression which are best fit for the following data:

х	1	2	3	4	5	6	7
Y	9	8	10	12	11	13	14

- 2. (a) The equations of two regression lines obtained in a correlation analysis are 6X + 10Y = 119 and 2X + 3Y = 12. The variance of Y is 4. Find
  - (i) The mean of X and Y
  - (ii) Coefficient of correlation
  - (iii) Variance of X
  - (iv) Covariance of X and Y

(b) Fit a least square curve  $y = ax^b$  to following data:

	<u>-</u>				
x	1	2	3	4	5
у	0.5	2	4.5	8	12.5

UNIT - II

- 3. (a) An urn contains 6 white, 4 rest and 9 black balls. If 3 balls are drawn at random, find the probability that (i) two of the balls are white, (ii) one of each colour (iii) none is red (iv) at least one is white.
  - (b) State and prove Boole's inequality.
- 4. (a) A letter is taken out at random out of "ASSISTANT" and a letter out of "STATISTIC". What is the chance that they are the same letters?
  - (b) Give the classical and statistical definitions of probability. What are the objections raised in these definitions?

UNIT - III

5. (a) The probabilities of X, Y, and Z becoming managers are  $\frac{4}{9}$ ,  $\frac{2}{9}$  and  $\frac{1}{3}$  respectively. The

(3)

P. T. O.

- probabilities that the Bonus scheme will be introduced if x, y and z becomes managers are  $\frac{3}{10}$ ,  $\frac{1}{2}$  and  $\frac{4}{5}$  respectively.
- (i) What is the probability that Bonus scheme will be introduce, and
- (ii) If the Bonus scheme has been introduced, what is the probability that manager appointed was X?

(b) A random variable X has the following probability distribution:

	IISUII	Duttor	<u> </u>					_	
x	0	1	2	3	4	5	6	7	8
p(x)	k	3k	5k	7k	9k	11k	13k	15k	17k

- (i) Determine the value of k.
- (ii) Find P(X < 3),  $P(X \ge 3)$ , P(0 < X < 5)
- (iii) What is the smallest value of x for which  $P(x \le x) > 0.5$ ?
- (iv) Find the distribution function of X.

6. (a) A r. v X has the density function:

$$f(x) = \begin{cases} \frac{1}{4}, & -2 < x < 2 \\ 0 & \text{elsewhere} \end{cases}$$

- (iii) P(2×+3>5 that the limit lain what is more quish between years (b) Explain what is meant by a random val random variable. Define distribution function random variable.

#### UNIT - IV

- 7. (a) The joint probability distribution of two rund variables X and Y is given by  $P(X = 0, Y \rightarrow 0)$  $=\frac{1}{3}$ ,  $P(x = 1, y = -1)\frac{1}{3}$ , p(X = 1, y = 1)Find
  - (i) Marginal distribution of X and Y and

(5)

- (ii) The conditional probability distribution of X given Y = 1.
- (b) Suppose that two dimensional continuous random variable (X, Y) has joint p. d. f given by:

$$f(x,y) = \begin{cases} 6x^2y, & 0 < x < 1, & 0 < y < 1 \\ 0 & \text{elsewhere} \end{cases}$$

- (i) Verify that  $\int_{0}^{1} \int_{0}^{1} f(x, y) dx dy = 1$
- (ii) Find  $P\left(0 < x < \frac{3}{4}, \frac{1}{3} < y < 2\right)$ P(X+Y<1), P(X>y)and P(X<1|Y<2)
- (a) What is the expectation of the number of failures preceding the first success in an infinite series of independent trails with constant probability p of saccess in each trial?
- (h) The p. d. f of a r. v is given by  $f(x) = kxe^{-x^2/2}$  $x \ge 0$ , when K is a constant. Evaluate K and obtain mean, median and mode of the distribution.



#### UNIT - V

- 9. (a) Distinguish between positive and man correlation.
  - (b) Fit a st line in the following data:

4	3	2	1	x
	-	2	2	.,

- (c) A, B, and C are three mutually exclusive at exhaustive events associated with a random experiment. Wind P(A) given that  $P(B) = \frac{3}{2}P(A)$  and  $P(C) = \frac{1}{2}P(B)$ .
- (d) State Baye's theorem
- (e) If  $p(x) = \begin{cases} \frac{x}{15}; & x = 1,2,3,4,5 \\ 0 & \text{elsewhere} \end{cases}$ Find  $\rho\left(\frac{1}{2} < x < \frac{5}{2} \mid x > 1\right)$
- (f) If  $\mu_1^1 = \frac{16}{7}$  and  $\mu_2^1 = \frac{40}{7}$ , find variance ( $\mu_2$ )