

94062

B. Sc. (Hons.) Mathematics
5th Semester Old/New Scheme
Examination – February, 2022

INTEGRAL EQUATION

Paper : BHM-354

Time : Three Hours] *Maximum Marks : 60*

Before answering the questions, candidates should ensure that they have been supplied the correct and complete question paper. No complaint in this regard, will be entertained after examination.

Note : Attempt five questions in all, selecting one question from each Section. Question No. 9 (Section-V) is compulsory. All questions carry equal marks.

SECTION – I

1. (a) Using the method of successive approximation solve the integral equation : 6

$$u(x) = x - \int_0^x (x - \xi) u(\xi) d\xi.$$

- (b) Solve the integral equation $\sin x = \lambda \int_0^x e^{x-\xi} u(\xi) d\xi$. 6

2. (a) By the Resolvent Kernel method, solve Volterra integral equation of second kind : 6

$$u(x) = f(x) + \lambda \int_0^x e^{x-\xi} u(\xi) d\xi.$$

- (b) Reduce the initial value problem $y'' + xy = 1$, $y'(0) = 0 = y(0)$ to Volterra integral equation. 6

SECTION – II

3. (a) Reduce the boundary value problem, 6
 $y'' + A(x)y' + B(x)y = g(x)$, $a \leq x \leq b$, $y(a) = c_1$, $y(b) = c_2$
 to a Fredholm integral equation.

- (b) Find the first two approximation of the solution of Fredholm integral equation : 6

$$u(x) = 1 + \int_0^1 K(x, \xi) u(\xi) d\xi \text{ where } K(x, \xi) = \begin{cases} x & 0 \leq x \leq \xi \\ \xi & \xi \leq x \leq 1 \end{cases}.$$

4. (a) Solve the following integral equation of second kind by the method of successive approximations to third order : 6

$$\phi(x) = 1 + \lambda \int_0^1 (x + \xi) \phi(\xi) d\xi, \phi_0(x) = 1.$$

- (b) Find the resolvent kernels for the Kernel $K(x, \xi) = e^{(x+\xi)}$, $a = 0$, $b = 1$. 6

SECTION - III

5. (a) Construct Green's function for the equation

$$x \frac{d^2 u}{dx^2} + \frac{du}{dx} = 0 \text{ with the conditions } u(x) \text{ is bounded}$$

as $x \rightarrow 0$, $u(1) = \mu u'(1)$, $\mu \neq 0$. 6

- (b) Reduce the Bessel's differential equation

$$x^2 \frac{d^2 u}{dx^2} + x \frac{du}{dx} + (\lambda x^2 - 1)u = 0 \text{ with the conditions}$$

$u(0) = 0$, $u(1) = 0$ into an integral equation, using Green's function. 6

6. (a) Reduce the boundary value problem

$$\frac{d^2 u}{dx^2} + \frac{\pi^2}{4} u = \lambda u + \cos \frac{\pi x}{2}, u(-1) = u(1) \text{ and } u'(-1) = u'(1)$$

to an integral equation, using Green's function. 6

- (b) Discuss construction of Green's function by variation of parameter method. 6

SECTION - IV

7. For the Fredholm integral equation $y(x) = \lambda \int_a^b K(x, \xi)$

$y(\xi) d\xi$ with symmetric Kernel, prove that : 12

- (i) The eigen functions corresponding to two different eigen values are orthogonal over (a, b) .
(ii) The eigen values are real.

8. (a) Determine the eigen values and eigen functions for the following integral equations : 6

$$\phi(x) = \lambda \int_0^1 (5x\xi^3 + 4x^2\xi)\phi(\xi) d\xi.$$

- (b) Solve the following integral equation by the method of Fredholm Resolvent Kernel : 6

$$\phi(x) = \left(\sin x - \frac{x}{4}\right) + \frac{1}{4} \int_0^{\pi/2} x\xi\phi(\xi) d\xi.$$

SECTION - V

9. (a) What is Fredholm integral equation of first kind and given an example ? 12

- (b) Show that the function $y(x) = (1+x^2)^{-\frac{3}{2}}$ is solution of the Volterra integral equation $y(x) = \frac{1}{1+x^2} -$

$$\int_0^x \frac{\xi}{1+x^2} y(\xi) d\xi.$$

- (c) What do you mean by initial value problem ?
(d) Write four basic properties of Green's function.
(e) What do you mean by homogenous integral equation of Fredholm type ?
(f) Define Iterated Kernel and Neumann series for Fredholm equation.