

Roll No. ....

41202

**B. Sc. (Pass Course) 4th Semester  
Examination – May, 2019**

**CHEMISTRY - II (Physical Chemistry)**

**Paper : CII-402**

*Time : Three hours ] [ Maximum Marks : 30*

*Before answering the questions, candidates should ensure that they have been supplied the correct and complete question paper. No complaint in this regard, will be entertained after examination.*

**Note :** Attempt *five* questions in all, including Question No. 1 which is *compulsory*. Select *one* question from each Section. All questions carry equal marks.

1. **Compulsory Questions :** 1,1, 1, 1, 1, 1

- (a) State second law of thermodynamics.
- (b) Define residual entropy.

P. T. O.

41202

- (c) What are reversible and irreversible cells ? Give examples.
- (d) Under what conditions  $\Delta G$  becomes equal to  $\Delta A$ .
- (e) What are the limitations of quinhydrone electrode ?
- (f) Why the efficiency of heat engine cannot be greater than one ?

**SECTION – A**

- 2. (a) Describe Carnot cycle and derive an expression for the efficiency of a heat engine working between temperatures  $T_1$  and  $T_2$ . 4
- (b) What is the criterion of spontaneity in terms of entropy ? 2
- 3. (a) Derive an expression for entropy as a function of  $V$  &  $T$  and  $P$  &  $T$  where  $V$ ,  $P$ ,  $T$  are volume, pressure and temperature respectively. 5
- (b) What is the criterion of spontaneity in terms of entropy ? 1

(2)

**SECTION – B**

4. (a) State third law of thermodynamics. How does it help in the determination of absolute entropies of chemical compounds at desired temperature? 2
- (b) The free energy change ( $\Delta G$ ) accompanying a given process is  $-85.77$  KJ at  $25^\circ\text{C}$  and  $-83.68$  KJ at  $35^\circ\text{C}$ . Calculate the change in enthalpy ( $\Delta H$ ) for the process at  $30^\circ\text{C}$ . 4
5. (a) Derive Gibb's Helmholtz equation. 5
- (b) What is the difference between Helmholtz function and Gibb's function? Under what conditions  $\Delta G$  becomes equal to  $\Delta A$ . 1

**SECTION – C**

6. (a) Derive the Nernst equation for EMF of a complete electrochemical cell. 4
- (b) Write a short note on single electrode potential. 2
7. (a) Calculate the EMF of a Zinc-silver cell at  $30^\circ\text{C}$  when the activity of  $\text{Zn}^{2+}$  ions is 0.5 and the activity of  $\text{Ag}^+$  ions is 10. Standard reduction potential at  $30^\circ\text{C}$  is (i)  $\text{Ag}^+/\text{Ag} = 0.799$  V (ii)  $\text{Zn}^{2+}/\text{Zn} = -0.76$  V. 4

(3)

P. T. O

- (b) Derive an expression for calculating mean ionic activity coefficient. 2

**SECTION – D**

8. Explain Concentration Cell (a) with transference (b) without transference. 3, 3
9. (a) Discuss the application of EMF measurement in potentiometric titrations. 4
- (b) Calculate the EMF of the following concentration cells at  $25^\circ\text{C}$ .  $\text{Ag}|0.001\text{N AgNO}_3||0.1\text{N AgNO}_3$ ; assuming the activities of silver ions to be equal to the concentrations. 2

(4)