FACULTY OF ENGINEERING

B.E. II-Semester (AICTE) (Main & Backlog) Examination, November 2020

Subject: Mathematics - II

Time: 2 Hours

Max. Marks: 70

Note: Answer Any five Questions from Part-A & Any Four Questions From Part-B.

PART - A (5x4=20 Marks)

- 1 Examine whether the vector (1, 2,), (3, 4), (3, 7) are linearly independent.
- 2 If 1, -1, 2 are the eigen values of a 3 x 3 matrix A, find the determinant of the matrix A³ 2A⁻¹ + I.
 - 3 Define exact differential equation.
- 4 Find the singular solution of the Clairant's equation y + y
- 5 Find the complementary function of $(D^2 + D + 1)^2y = e^{-t}$
- 6 Solve $x \frac{d^2 y}{dx^2} + \frac{dy}{dx} = 0$.
- 7 Evaluate $r\left(-\frac{3}{2}\right)$.
- 8 State Rodrigue's formula and hence find P2
- 9 Find L{e-! sint cost}
- 10 Evaluate $\int_{t}^{\sin t} dt$ using Laplate transform.

FRT - B (4x15=60 Marks)

11 (a) Test for consiste and hence solve the following system of equations.

$$x_1 + 2x_2 + x_3 = 2$$
 $3x_1 + x_2 - 2x_1 = 1$, $4x_1 - 3x_2 - x_1 = 3$, $2x_1 + 4x_2 + 2x_3 = 4$

- (b) Find the characteristics equation of $A = \begin{pmatrix} 4 & 3 & 1 \\ 2 & 1 & -2 \\ 1 & 2 & 1 \end{pmatrix}$ and hence find A⁻¹.
- 12 (a) Solve $(3x^2y^4 + 2xy)dx + (2x^3y^3 x^2)dy = 0$.p
 - (b) Find the orthogonal trajectories of the family of parabolas $y^2 = 2cx + c^2$.
- 13 (a) Find the general solution of the differential equation

$$\frac{d^3y}{d^3} - y = (e^x + 1)^2.$$

- (b) Solve $y'' + 2y' + 2y = e^{-x} \cos_x by$ the method of variation of parameters.
- 14 (a) Evaluate $\int_{a}^{1} \frac{dx}{\sqrt{1-x^4}}$ using Beta and Gamma functions.
 - (b) Show that $P_{2n}(0) = (-1)^n \frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n-1)}{2 \cdot 4 \cdot 6 \cdot \dots \cdot 2n}$ and $P_{2n+1}(0)=0$.

- 15 (a) Find the inverse Laplace transform of $\log \left(\frac{5+a}{5+b} \right)$.
 - (b) Apply Laplace transforms to solve $y'' + y = 3\cos 2x$, y'(0) = 0 = y(0).
- 46 Reduce the quadratic form Q = 2(xy + yz + zx) to Canonical form using orthogonal transformation.
- 17 (a) Show that $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$.
 - (b) Apply convolution theorem to find $L^{-1}\left\{\frac{s}{(s^2+1)(s-1)}\right\}$